

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C07C 311/29, C07D 295/18, 213/56, A61K 31/535, 31/44, 31/495, 31/40, 31/18		A1	(11) International Publication Number: WO 96/27583
			(43) International Publication Date: 12 September 1996 (12.09.96)
(21) International Application Number: PCT/US96/02679 (22) International Filing Date: 7 March 1996 (07.03.96) (30) Priority Data: 08/401,049 8 March 1995 (08.03.95) US (60) Parent Application or Grant (63) Related by Continuation US 08/401,049 (CON) Filed on 8 March 1995 (08.03.95) (71) Applicant (for all designated States except US): PFIZER INC. [US/US]; 235 East 42nd Street, New York, NY 10017-5755 (US). (72) Inventors; and (75) Inventors/Applicants (for US only): ROBINSON, Ralph, P. [US/US]; 30 Friartuck Drive, Gales Ferry, CT 06335 (US). RIZZI, James, P. [US/US]; 34 Devonshire Drive, Waterford, CT 06335 (US).		(81) Designated States: AL, AM, AT, AU, AZ, BB, BG, BR, BY, CA, CH, CN, CZ, DE, DK, EE, ES, FI, GB, GE, HU, IS, JP, KE, KG, KP, KR, KZ, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, TJ, TM, TR, TT, UA, UG, US, UZ, VN, ARIPO patent (KE, LS, MW, SD, SZ, UG), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG). Published With international search report.	
(54) Title: ARYLSULFONYLAMINO HYDROXAMIC ACID DERIVATIVES (57) Abstract A compound of formula (I), wherein n, X, R ³ , R ⁴ and Ar are as defined above, useful in the treatment of a condition selected from the group consisting of arthritis, cancer, tissue ulceration, restenosis, periodontal disease, epidermolysis bullosa, scleritis and other diseases characterized by matrix metalloproteinase activity, AIDS, sepsis, septic shock and other diseases involving the production of TNF.			
<div style="text-align: center;"> <p style="text-align: right;">(I)</p> </div>			

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AM	Armenia	GB	United Kingdom	MW	Malawi
AT	Austria	GE	Georgia	MX	Mexico
AU	Australia	GN	Guinea	NE	Niger
BB	Barbados	GR	Greece	NL	Netherlands
BE	Belgium	HU	Hungary	NO	Norway
BF	Burkina Faso	IE	Ireland	NZ	New Zealand
BG	Bulgaria	IT	Italy	PL	Poland
BJ	Benin	JP	Japan	PT	Portugal
BR	Brazil	KE	Kenya	RO	Romania
BY	Belarus	KG	Kyrgyzstan	RU	Russian Federation
CA	Canada	KP	Democratic People's Republic of Korea	SD	Sudan
CF	Central African Republic	KR	Republic of Korea	SE	Sweden
CG	Congo	KZ	Kazakhstan	SG	Singapore
CH	Switzerland	LI	Liechtenstein	SI	Slovenia
CI	Côte d'Ivoire	LK	Sri Lanka	SK	Slovakia
CM	Cameroon	LR	Liberia	SN	Senegal
CN	China	LT	Lithuania	SZ	Swaziland
CS	Czechoslovakia	LU	Luxembourg	TD	Chad
CZ	Czech Republic	LV	Latvia	TG	Togo
DE	Germany	MC	Monaco	TJ	Tajikistan
DK	Denmark	MD	Republic of Moldova	TT	Trinidad and Tobago
EE	Estonia	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	UG	Uganda
FI	Finland	MN	Mongolia	US	United States of America
FR	France	MR	Mauritania	UZ	Uzbekistan
GA	Gabon			VN	Viet Nam

- 1 -

ARYLSULFONYLAMINO HYDROXAMIC ACID DERIVATIVESBackground of the Invention

The present invention relates to arylsulfonylamino hydroxamic acid derivatives which are
5 inhibitors of matrix metalloproteinases or the production of tumor necrosis factor (hereinafter also referred to as TNF) and as such are useful in the treatment of a condition selected from the group consisting of arthritis, cancer, tissue ulceration,
10 restenosis, periodontal disease, epidermolysis bullosa, scleritis and other diseases characterized by matrix metalloproteinase activity, AIDS, sepsis, septic shock and other diseases involving the production of TNF.

This invention also relates to a method of
15 using such compounds in the treatment of the above diseases in mammals, especially humans, and to the pharmaceutical compositions useful therefor.

There are a number of enzymes which effect the breakdown of structural proteins and which are
20 structurally related metalloproteases. Matrix-degrading metalloproteinases, such as gelatinase, stromelysin and collagenase, are involved in tissue matrix degradation (e.g. collagen collapse) and have been implicated in many pathological conditions
25 involving abnormal connective tissue and basement membrane matrix metabolism, such as arthritis (e.g. osteoarthritis and rheumatoid arthritis), tissue ulceration (e.g. corneal, epidermal and gastric ulceration), abnormal wound healing, periodontal

- 2 -


diseases, bone diseases (e.g. Paget's disease and osteoporosis), tumor metastasis or invasion, as well as HIV-infection (J. Leuk. Biol., 52 (2):244-248, 1992).

5 Tumor necrosis factor is recognized to be involved in many infectious and auto-immune diseases (W. Friers, FEBS Letters, 1991, 285, 199).

Furthermore, it has been shown that TNF is the prime mediator of the inflammatory response seen in sepsis and septic shock (C.E. Spooner et al., Clinical Immunology and Immunopathology, 1992, 62 S11).

Summary of the Invention

The present invention relates to a compound of the formula



or the pharmaceutically acceptable salts thereof, wherein

25 n is 1 to 6;

X is hydroxy, (C₁-C₆)alkoxy or NR¹R² wherein R¹ and R² are each independently selected from the group consisting of hydrogen, (C₁-C₆)alkyl, piperidyl, (C₁-C₆)alkylpiperidyl, (C₆-C₁₀)aryl piperidyl, (C₅-C₉)heteroaryl piperidyl, (C₆-C₁₀)aryl (C₁-C₆)alkylpiperidyl, (C₅-C₉)heteroaryl (C₁-C₆)alkylpiperidyl, (C₁-C₆)acylpiperidyl, (C₆-C₁₀)aryl, (C₅-C₉)heteroaryl, (C₆-C₁₀)aryl (C₁-C₆)alkyl, (C₅-C₉)heteroaryl (C₁-C₆)alkyl, (C₆-C₁₀)aryl (C₆-C₁₀)aryl, (C₆-C₁₀)aryl (C₆-C₁₀)aryl (C₁-C₆)alkyl, (C₃-C₆)cycloalkyl, (C₃-C₆)cycl alkyl (C₁-C₆)alkyl, R⁵(C₂-

30


35

- 3 -

- C_6)alkyl, (C_1-C_5) alkyl(CHR^5) (C_1-C_6) alkyl where in R^5 is hydroxy, (C_1-C_6) acyloxy, (C_1-C_6) alkoxy, piperazino, (C_1-C_6) alkylamino, (C_1-C_6) alkylthio, (C_6-C_{10}) arylthio, (C_1-C_6) alkylsulfinyl, (C_6-C_{10}) arylsulfinyl, (C_1-C_6) alkylsulfoxyl, (C_6-C_{10}) arylsulfoxyl, amino, (C_1-C_6) alkylamino, $((C_1-C_6)alkyl)_2$ amino, (C_1-C_6) acylpiperazino, (C_1-C_6) alkylpiperazino, (C_6-C_{10}) aryl (C_1-C_6) alkylpiperazino, (C_5-C_9) heteroaryl (C_1-C_6) alkylpiperazino, morpholino, thiomorpholino,
- 10 piperidino or pyrrolidino; R^6 (C_1-C_6) alkyl, (C_1-C_5) alkyl(CHR^6) (C_1-C_6) alkyl wherein R^6 is piperidyl, (C_1-C_6) alkylpiperidyl, (C_6-C_{10}) aryl piperidyl, (C_6-C_{10}) aryl (C_1-C_6) alkylpiperidyl, (C_5-C_9) heteroaryl piperidyl or (C_5-C_9) heteroaryl (C_1-C_6) alkylpiperidyl; and $CH(R^7)COR^8$ wherein R^7 is
- 15 hydrogen, (C_1-C_6) alkyl, (C_6-C_{10}) aryl (C_1-C_6) alkyl, (C_5-C_9) heteroaryl (C_1-C_6) alkyl, (C_1-C_6) alkylthio (C_1-C_6) alkyl, (C_6-C_{10}) arylthio (C_1-C_6) alkyl, (C_1-C_6) alkylsulfinyl (C_1-C_6) alkyl, (C_6-C_{10}) arylsulfinyl (C_1-C_6) alkyl, (C_1-C_6) alkylsulfonyl (C_1-C_6) alkyl, (C_6-C_{10}) arylsulfonyl (C_1-C_6) alkyl, hydroxy (C_1-C_6) alkyl, amino (C_1-C_6) alkyl, (C_1-C_6) alkylamino (C_1-C_6) alkyl, $((C_1-C_6)$ alkylamino) $_2$ (C_1-C_6) alkyl, $R^9R^{10}NCO(C_1-C_6)$ alkyl or $R^9OCO(C_1-C_6)$ alkyl wherein R^9 and R^{10} are each
- 20 independently selected from the group consisting of hydrogen, (C_1-C_6) alkyl, (C_6-C_{10}) aryl (C_1-C_6) alkyl and (C_5-C_9) heteroaryl (C_1-C_6) alkyl; and R^8 is $R^{11}O$ or $R^{11}R^{12}N$ wherein R^{11} and R^{12} are each independently selected from the group consisting of hydrogen, (C_1-C_6) alkyl, (C_6-C_{10}) aryl (C_1-C_6) alkyl and (C_5-C_9) heteroaryl (C_1-C_6) alkyl;
- 30 or R^1 and R^2 , or R^9 and R^{10} , or R^{11} and R^{12} may be taken together to form an azetidiny, pyrrolidinyl, morpholinyl, thiomorpholinyl, indolinyl,
- 35 isoindolinyl, tetrahydroquinolinyl, tetrahydr isoquinolinyl, (C_1-C_6) acylpiperazinyl, (C_1-C_6) alkylpiperazinyl, (C_6-C_{10}) aryl piperazinyl, (C_5-C_9) heteroaryl piperazinyl,

- 4 -

C₉)heteroarylpiperazinyl or a bridged diazabicycloalkyl ring selected from the group consisting of



wherein r is 1, 2 or 3;

m is 1 or 2;

20 p is 0 or 1; and

Q is hydrogen, (C₁-C₃)alkyl or (C₁-C₆)acyl;

R³ and R⁴ are each independently selected

from the group consisting of hydrogen, (C₁-C₆)alkyl,

trifluoromethyl, trifluoromethyl(C₁-C₆)alkyl, (C₁-

25 C₆)alkyl (difluoromethylene), (C₁-

C₃)alkyl(difluoromethylene)(C₁-C₃)alkyl, (C₆-C₁₀)aryl,

(C₅-C₉)heteroaryl, (C₆-C₁₀)aryl(C₁-C₆)alkyl, (C₅-

C₉)heteroaryl(C₁-C₆)alkyl, (C₆-C₁₀)aryl(C₆-C₁₀)aryl,

(C₆-C₁₀)aryl(C₆-C₁₀)aryl(C₁-C₆)alkyl, (C₃-


30 C₆)cycloalkyl, (C₃-C₆)cycloalkyl(C₁-C₆)alkyl,

hydroxy(C₁-C₆)alkyl, (C₁-C₆)acyloxy(C₁-C₆)alkyl, (C₁-

C₆)alkoxy(C₁-C₆)alkyl, piperazinyl(C₁-C₆)alkyl, (C₁-

- 5 -

- C_6)acylamino(C_1-C_6)alkyl, piperidyl, (C_1-C_6)alkylpiperidyl, (C_6-C_{10})aryl(C_1-C_6)alkoxy(C_1-C_6)alkyl, (C_5-C_9)heteroaryl(C_1-C_6)alkoxy(C_1-C_6)alkyl, (C_1-C_6)alkylthio(C_1-C_6)alkyl, (C_6-C_{10})arylthio(C_1-C_6)alkyl, (C_1-C_6)alkylsulfinyl(C_1-C_6)alkyl, (C_6-C_{10})arylsulfinyl(C_1-C_6)alkyl, (C_1-C_6)alkylsulfonyl(C_1-C_6)alkyl, (C_6-C_{10})arylsulfonyl(C_1-C_6)alkyl, amino(C_1-C_6)alkyl, (C_1-C_6)acylpiperazino, (C_5-C_9)heteroarylpiperazino, (C_1-C_6)alkylpiperazino, (C_6-C_{10})aryl(C_1-C_6)alkylpiperazino, (C_5-C_9)heteroaryl(C_1-C_6)alkylpiperazino, morpholino, thiomorpholino, piperidino, pyrrolidino, piperidyl, (C_1-C_6)alkylpiperidyl, (C_6-C_{10})aryl(C_1-C_6)alkylpiperidyl, (C_5-C_9)heteroarylpiperidyl, (C_6-C_{10})aryl(C_1-C_6)alkylpiperidyl, (C_5-C_9)heteroaryl(C_1-C_6)alkylpiperidyl or (C_1-C_6)acylpiperidyl;
- or R^3 and R^4 , or R^{20} and R^{21} may be taken together to form a (C_3-C_6)cycloalkyl, oxacyclohexyl, thiocyclohexyl, indanyl or tetralinyl ring or a group of the formula



- wherein R^{15} is hydrogen, (C_1-C_6)acyl, (C_1-C_6)alkyl, (C_6-C_{10})aryl(C_1-C_6)alkyl, (C_5-C_9)heteroaryl(C_1-C_6)alkyl or (C_1-C_6)alkylsulfonyl; and
- Ar is (C_6-C_{10})aryl, (C_5-C_9)heteroaryl, (C_1-C_6)alkyl(C_6-C_{10})aryl, (C_1-C_6)alkoxy(C_6-C_{10})aryl, ((C_1-C_6)alkylamino) $_2$ (C_1-C_6)alkyl, $R^{13}CO(C_1-C_6)$ alkyl wherein R^{13} is $R^{20}O$ or $R^{20}R^{21}N$ wherein R^{20} and R^{21} are each independently selected from the group consisting of hydrogen, (C_1-C_6)alkyl, (C_6-C_{10})aryl(C_1-C_6)alkyl or (C_5-C_9)heteroaryl(C_1-C_6)alkyl; or $R^{14}(C_1-C_6)$ alkyl wherein R^{14} is (C_1-C_6)acylpiperazino, (C_6-C_{10})arylpiperazino, (C_5-C_9)heteroarylpiperazino, (C_1-C_6)alkylpiperazino, (C_6-C_{10})aryl(C_1-C_6)alkylpiperazino, (C_5-C_9)heteroaryl(C_1-C_6)alkylpiperazino, morpholino, thiomorpholino, piperidino, pyrrolidino, piperidyl, (C_1-C_6)alkylpiperidyl, (C_6-C_{10})aryl(C_1-C_6)alkylpiperidyl, (C_5-C_9)heteroarylpiperidyl, (C_6-C_{10})aryl(C_1-C_6)alkylpiperidyl, (C_5-C_9)heteroaryl(C_1-C_6)alkylpiperidyl or (C_1-C_6)acylpiperidyl;

- 6 -

C₆)alkoxy)₂(C₆-C₁₀)aryl, (C₆-C₁₀)aryloxy(C₆-C₁₀)aryl,
(C₅-C₉)heteroaryloxy(C₆-C₁₀)aryl, (C₁-C₆)alkyl(C₅-
C₉)heteroaryl, (C₁-C₆)alkoxy(C₅-C₉)heteroaryl, ((C₁-
C₆)alkoxy)₂(C₅-C₉)heteroaryl, (C₆-C₁₀)aryloxy(C₅-
5 C₉)heteroaryl, (C₅-C₉)heteroaryloxy(C₅-C₉)heteroaryl;
with the proviso that when either R¹ or R² is
CH(R⁷)COR⁸ wherein R⁷ and R⁸ are as defined above, the
other of R¹ or R² is hydrogen, (C₁-C₆)alkyl or benzyl.

The term "alkyl", as used herein, unless
10 otherwise indicated, includes saturated monovalent
hydrocarbon radicals having straight, branched or
cyclic moieties or combinations thereof.

The term "alkoxy", as used herein, includes
O-alkyl groups wherein "alkyl" is defined above.

15 The term "aryl", as used herein, unless
otherwise indicated, includes an organic radical
derived from an aromatic hydrocarbon by removal of one
hydrogen, such as phenyl or naphthyl, optionally
substituted by 1 to 3 substituents selected from the
20 group consisting of fluoro, chloro, trifluoromethyl,
(C₁-C₆)alkoxy, (C₆-C₁₀)aryloxy, trifluoromethoxy,
difluoromethoxy and (C₁-C₆)alkyl.

The term "heteroaryl", as used herein, unless
otherwise indicated, includes an organic radical
25 derived from an aromatic heterocyclic compound by
removal of one hydrogen, such as pyridyl, furyl,
pyrolyl, thienyl, isothiazolyl, imidazolyl,
benzimidazolyl, terazolyl, pyrazinyl, pyrimidyl,
quinolyl, isoquinolyl, benzofuryl, isobenzofuryl,
30 benzothienyl, pyrazolyl, indolyl, isoindolyl, purinyl,
carbazolyl, isoxazolyl, thiazolyl, oxazolyl,
benzthiazolyl or benzoxazolyl, optionally substituted
by 1 to 2 substituents selected from the group
consisting of fluoro, chloro, trifluoromethyl, (C₁-
35 C₆)alkoxy, (C₆-C₁₀)aryloxy, trifluoromethoxy,
difluoromethoxy and (C₁-C₆)alkyl.

The term "acyl", as used herein, unless

- 7 -

otherwise indicated, includes a radical of the general formula RCO wherein R is alkyl, alkoxy, aryl, arylalkyl or aryalkyloxy and the terms "alkyl" or "aryl" are as defined above.

5 The term "acyloxy", as used herein, includes O-acyl groups wherein "acyl" is defined above.

 The compound of formula I may have chiral centers and therefore exist in different enantiomeric forms. This invention relates to all optical isomers
10 and stereoisomers of the compounds of formula I and mixtures thereof.

 Preferred compounds of formula I include those wherein n is 2.

 Other preferred compounds of formula I
15 include those wherein Ar is 4-methoxyphenyl or 4-phenoxyphenyl.

 Other preferred compounds of formula I include those wherein either R³ or R⁴ is not hydrogen.


 Other preferred compounds of formula I
20 include those wherein n is 1 and either R¹ or R² is hydrogen.

 Other preferred compounds of formula I include those wherein X is hydroxy, Ar is 4-methoxyphenyl or 4-phenoxyphenyl and either R³ or R⁴ is
25 not hydrogen.

 Other preferred compounds of formula I include those wherein X is alkoxy, Ar is 4-methoxyphenyl or 4-phenoxyphenyl and either R³ or R⁴ is not hydrogen.

30 Other preferred compounds of formula I include those wherein Ar is 4-methoxyphenyl or 4-phenoxyphenyl and R³ and R⁴ are taken together to form (C₃-C₆)cycloalkanyl, oxacyclohexanyl, thiocyclohexanyl, indanyl or a group of the formula

- 8 -



5

wherein R¹⁵ is (C₁-C₆)acyl, (C₁-C₆)alkyl, (C₆-C₁₀)aryl(C₁-C₆)alkyl, (C₅-C₉)heteroaryl(C₁-C₆)alkyl or (C₁-C₆)alkylsulfonyl.

10

More preferred compounds of formula I are those wherein n is 2, Ar is 4-methoxyphenyl or 4-phenoxyphenyl, R¹ and R² are taken together to form piperazinyl, (C₁-C₆)alkylpiperazinyl, (C₆-C₁₀)aryl piperazinyl or (C₅-C₉)heteroaryl(C₁-

15

C₆)alkylpiperazinyl, and either R³ or R⁴ is not hydrogen or both R³ and R⁴ are not hydrogen.

More preferred compounds of formula I are those wherein n is 2, Ar is 4-methoxyphenyl or 4-phenoxyphenyl, R¹ is hydrogen or (C₁-C₆)alkyl, R² is 2-pyridylmethyl, 3-pyridylmethyl or 4-pyridylmethyl, and either R³ or R⁴ is not hydrogen or both R³ and R⁴ are not hydrogen.

20

More preferred compounds of formula I are those wherein n is 1, Ar is 4-methoxyphenyl or 4-phenoxyphenyl, R¹ is hydrogen, R² is 2-pyridylmethyl, 3-pyridylmethyl or 4-pyridylmethyl, and either R³ or R⁴ is not hydrogen or both R³ and R⁴ are not hydrogen.

25

More preferred compounds of formula I are those wherein n is 2, Ar is 4-methoxyphenyl, R¹ is hydrogen or (C₁-C₆)alkyl and R² is R⁵ (C₂-C₆)alkyl wherein R⁵ is morpholino, thiomorpholino, piperidino, pyrrolidino, (C₁-C₆)acylpiperazino, (C₁-C₆)alkylpiperazino, (C₆-C₁₀)arylpiperazino, (C₅-C₉)heteroarylpiperazino, (C₆-C₁₀)aryl(C₁-C₆)alkylpiperazino or (C₅-C₉)heteroaryl(C₁-C₆)alkylpiperazino and either R³ or R⁴ is not hydrogen or both R³ and R⁴ are not hydrogen.

30

35

More preferred compounds of formula I are

- 9 -

those wherein n is 1, Ar is 4-methoxyphenyl or 4-phenoxyphenyl, R¹ is hydrogen, R² is R⁵ (C₂-C₆)alkyl wherein R⁵ is morpholino, thiomorpholino, piperidino, pyrrolidino (C₁-C₆)acylpiperazino, (C₁-C₆)alkylpiperazino, (C₆-C₁₀)arylpiperazino, (C₅-C₉)heteroarylpiperazino, (C₆-C₁₀)aryl(C₁-C₆)alkylpiperazino or (C₅-C₉)heteroaryl(C₁-C₆)alkylpiperazino and either R³ or R⁴ is not hydrogen or both R³ and R⁴ are not hydrogen.

Specific preferred compounds of formula I include the following:

2-(R)-N-Hydroxy-2-[(4-methoxybenzenesulfonyl)(3-morpholin-4-yl-3-oxopropyl)amino]-3-methylbutyramide;

2-(R)-2-[(2-Benzylcarbamoyl)ethyl](4-methoxybenzenesulfonyl)amino]-N-hydroxy-3-methylbutyramide;

2-(R)-N-Hydroxy-2-[(4-methoxybenzenesulfonyl)(2-[(pyridin-3-ylmethyl)carbamoyl]ethyl)amino]-3-methylbutyramide;

2-(R)-N-Hydroxy-2-[(4-methoxybenzenesulfonyl)[2-(methylpyridin-3-ylmethylcarbamoyl)ethyl]amino]-3-methylbutyramide;

4-(3-[1-(R)-1-Hydroxycarbamoyl-2-methylpropyl](4-methoxybenzenesulfonyl)amino)propionyl)piperazine-1-carboxylic acid, tert-butyl ester;

2-(R)-N-Hydroxy-2-[(4-methoxybenzenesulfonyl)(3-oxo-3-piperazin-1-ylpropyl)amino]-3-methylbutyramide hydrochloride;

2-(R)-2-[(Benzylcarbamoylmethyl)(4-methoxybenzenesulfonyl)amino]N-hydroxy-3-methylbutyramide;

2-(R)-N-Hydroxy-2-[(4-methoxybenzenesulfonyl)-[(2-morpholin-4-ylethylcarbamoyl)methyl]amino]-3-methylbutyramide; and

2-(R)-N-Hydroxy-2-[(4-

- 10 -

methoxybenzenesulfonyl) [(pyridin-3-ylmethyl) carbamoyl]methyl) amino) -3-methylbutyramide.

Other specific compounds of formula I include the following:

- 5 2 - (R) - 3,3,3-Trifluoro-N-hydroxy-2-
[(methoxybenzenesulfonyl) (3-morpholin-4-yl-3-oxopropyl) amino] propionamide;
- 2 - (R) - N-Hydroxy-2 - ((4-phenoxybenzenesulfonyl) [2 - (methylpyridin-4-ylmethylcarbamoyl) ether] amino) -3-methylbutyramide;
- 10 4 - [4-Methoxybenzenesulfonyl) (3-morpholin-4-yl-3-oxopropyl) amino] -1-methylpiperidene-4-carboxylic acid hydroxyamide;
- 2 - (R) - N-Hydroxy-2 - ((4-methoxybenzenesulfonyl) - [3 - (4-methylpiperazin-1-yl) -3-oxopropyl] amino) -3-methylbutyramide;
- 15 2 - (R) - 2 - [(2-Carboxyethyl) (4-methoxybenzenesulfonyl) amino] -N-hydroxy-3-methylbutyramide;
- 20 [(2-Carboxyethyl) (3,4-dimethoxybenzenesulfonyl) amino] -N-hydroxy-acetamide;
- 2 - (R) - 2 - [(2-Carbamoyl ethyl) (4-methoxybenzenesulfonyl) amino] -N-hydroxy-3-methylbutyramide;
- 25 2 - (R) , 3 - (R) - 3, N-Dihydroxy-2 - [(4-methoxybenzenesulfonyl) (3-oxo-3-piperidin-1-ylpropyl) amino] -butyramide;
- 2 - (R) - N-Hydroxy-2 - ((4-methoxybenzenesulfonyl) [3 - (methylpyridin-3-ylmethylcarbamoyl) propyl] amino) -3-methylbutyramide;
- 30 2 - (R) - N-Hydroxy-2 - ((4-methoxybenzenesulfonyl) [2 - (methylcarboxymethylcarbamoyl) ethyl] amino) -3-methylbutyramide;
- 35 2 - (R) - N-Hydroxy-2 - ((4-methoxybenzenesulfonyl) - [(1-methylpiperidin-4-ylcarbamoyl) methyl] amino) -3-methylbutyramide;

- 11 -

2-(R)-2-Cyclohexyl-N-hydroxy-2-((4-methoxybenzenesulfonyl)-[3-(4-methylpiperazin-1-yl)-3-oxopropyl]amino)-acetamide;

2-(R)-N-Hydroxy-2-

- 5 [(methoxybenzenesulfonyl)(3-morpholin-4-yl-3-oxopropyl)amino]-4-(morpholin-4-yl)butyramide.

The present invention also relates to a pharmaceutical composition for (a) the treatment of a condition selected from the group consisting of

10 arthritis, cancer, tissue ulceration, restenosis, periodontal disease, epidermolysis bullosa, scleritis and other diseases characterized by matrix metalloproteinase activity, AIDS, sepsis, septic shock and other diseases involving the production of tumor

15 necrosis factor (TNF) or (b) the inhibition of matrix metalloproteinases or the production of tumor necrosis factor (TNF) in a mammal, including a human, comprising an amount of a compound of claim 1 or a pharmaceutically acceptable salt thereof, effective in

20 such treatments and a pharmaceutically acceptable carrier.

The present invention also relates to a method for the inhibition of (a) matrix metalloproteinases or (b) the production of tumor

25 necrosis factor (TNF) in a mammal, including a human, comprising administering to said mammal an effective amount of a compound of claim 1 or a pharmaceutically acceptable salt thereof.

The present invention also relates to a

30 method for treating a condition selected from the group consisting of arthritis, cancer, tissue ulceration, restenosis, periodontal disease, epidermolysis bullosa, scleritis and other diseases characterized by matrix metalloproteinase activity, AIDS, sepsis, septic shock

35 and other diseases involving the production of tumor necrosis factor (TNF) in a mammal, including a human, comprising administering to said mammal an amount of a


- 12 -

compound of claim 1 or a pharmaceutically acceptable salt thereof, effective in treating such a condition.


Detailed Description of the Invention

5 The following reaction Schemes illustrate the preparation of the compounds of the present invention. Unless otherwise indicated R^1 , R^2 , R^3 , R^4 , n and Ar in the reaction Schemes and the discussion that follow are defined as above.


- 13 -

Scheme 1


- 14 -

Scheme 1 cont'd


- 15 -

Scheme 2

- 16 -

Scheme 3

- 18 -

Scheme 4 cont'd

- 19 -

In reaction 1 of Scheme 1, the amino acid compound of formula VII, wherein R^{16} is (C_1-C_6) alkyl, benzyl, allyl or tert-butyl, is converted to the corresponding compound of formula VI by reacting VII with a reactive functional derivative of an arylsulfonic acid compound, such as an arylsulfonyl chloride, in the presence of a base, such as triethylamine, and a polar solvent, such as tetrahydrofuran, dioxane, water or acetonitrile, preferably a mixture of dioxane and water. The reaction mixture is stirred, at room temperature, for time period between about 10 minutes to about 24 hours, preferably about 60 minutes.

In reaction 2 of Scheme 1, the arylsulfonyl amino compound of formula VI, wherein R^{16} is (C_1-C_6) alkyl, benzyl, allyl or tert-butyl, is converted to the corresponding compound of formula V, wherein n is 1, 3, 4, 5 or 6, by reacting VI with a reactive derivative of an alcohol of the formula



such as the chloride, bromide or iodide derivative, preferably the bromide derivative, wherein the R^{17} protecting group is (C_1-C_6) alkyl, benzyl, allyl or tert-butyl, in the presence of a base such as potassium carbonate or sodium hydride, preferably sodium hydride, and a polar solvent, such as dimethylformamide. The reaction mixture is stirred, at room temperature, for a time period between about 60 minutes to about 48 hours, preferably about 18 hours. The R^{17} protecting group is chosen such that it may be selectively removed in the presence of and without loss of the R^{16} protecting group, therefore, R^{17} cannot be the same as R^{16} . Removal of the R^{17} protecting group from the compound of formula V to give the corresponding carboxylic acid of formula IV, in reaction 3 of Scheme 1, is carried out under conditions appropriate for that particular

- 20 -

R^{17} protecting group in use which will not affect the R^{16} protecting group. Such conditions include; (a) saponification where R^{17} is (C_1-C_6) alkyl and R^{16} is tert-butyl, (b) hydrogenolysis where R^{17} is benzyl and R^{16} is tert-butyl or (C_1-C_6) alkyl, (c) treatment with a strong acid such as trifluoroacetic acid or hydrochloric acid where R^{17} is tert-butyl and R^{16} is (C_1-C_6) alkyl, benzyl or allyl, or (d) treatment with tributyltinhydride and acetic acid in the presence of catalytic bis(triphenylphosphine) palladium (II) chloride where R^{17} is allyl and R^{16} is (C_1-C_6) alkyl, benzyl or tert-butyl.

In reaction 4 of Scheme 1, the carboxylic acid of formula IV is condensed with an amine, R^1R^2NH , or the salt thereof, to give the corresponding amide compound of formula III. The formation of amides from primary or secondary amines or ammonia and carboxylic acids is achieved by conversion of the carboxylic acid to an activated functional derivative which subsequently undergoes reaction with a primary or secondary amine or ammonia to form the amide. The activated functional derivative may be isolated prior to reaction with the primary or secondary amine or ammonia. Alternatively, the carboxylic acid may be treated with oxalyl chloride or thionyl chloride, neat or in an inert solvent, such as chloroform, at a temperature between about 25°C. to about 80°C., preferably about 50°C., to give the corresponding acid chloride functional derivative. The inert solvent and any remaining oxalyl chloride or thionyl chloride is then removed by evaporation under vacuum. The remaining acid chloride functional derivative is then reacted with the primary or secondary amine or ammonia in an inert solvent, such as methylene chloride, to form the amide. The preferred method for the condensation of the carboxylic acid of formula IV with an amine to provide the corresponding amide compound of

- 21 -

formula III is the treatment of IV with (benzotriazol-1-yloxy)tris(dimethylamin) phosphonium hexafluorophosphate in the presence of a base, such as triethylamine, to provide the benzotriazol-1-oxy ester
5 in situ which, in turn, reacts with the amine, R^1R^2N , in an inert solvent, such as methylene chloride, at room temperature to give the amide compound of formula III.

Removal of the R^{16} protecting group from the
10 compound of formula III to give the corresponding carboxylic acid of formula II, in reaction 5 of Scheme 1, is carried out under conditions appropriate for the particular R^{16} protecting group in use. Such conditions include; (a) saponification where R^{16} is
15 lower alkyl, (b) hydrogenolysis where R^{16} is benzyl, (c) treatment with a strong acid, such as trifluoroacetic acid or hydrochloric acid, where R^{16} is tert-butyl, or (d) treatment with tributyltinhydride and acetic acid in the presence of catalytic
20 bis(triphenylphosphine) palladium (II) chloride where R^{16} is allyl.

In reaction 6 of Scheme 1, the carboxylic acid compound of formula II is converted to the hydroxamic acid compound of formula I by treating II
25 with 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide and 1-hydroxybenztriazole in a polar solvent, such as dimethylformamide, followed by the addition of hydroxylamine to the reaction mixture after a time period between about 15 minutes to about 1 hour,
30 preferably about 30 minutes. The hydroxylamine is preferably generated in situ from a salt form, such as hydroxylamine hydrochloride, in the presence of a base, such as N-methylmorpholine. Alternatively, a protected derivative of hydroxylamine or its salt form, where the
35 hydroxyl group is protected as a tert-butyl, benzyl or allyl ether, may be used in the presence of (benz triazol-1-yloxy)tris(dimethylamino) phosphonium

- 22 -

hexafluor phosphate and a base, such as N-methylmorpholine. Removal of the hydroxylamine protecting group is carried out by hydrogenolysis for a benzyl protecting group or treatment with a strong acid, such as trifluoroacetic acid, for a tert-butyl protecting group. The allyl protecting group may be removed by treatment with tributyltinhydride and acetic acid in the presence of catalytic bis(triphenylphosphine) palladium (II) chloride. N,O-bis(4-methoxybenzyl)hydroxylamine may also be used as the protected hydroxylamine derivative where deprotection is achieved using a mixture of methanesulfonic acid and trifluoroacetic acid.

In reaction 1 of Scheme 2, the arylsulfonylamino compound of formula VI, wherein R^{16} is (C_1-C_6) alkyl, benzyl or tert-butyl, is converted to the corresponding compound of formula VIII, wherein R^{18} is 2-propenyl or 3-butenyl, by reacting IX with a reactive functional derivative, such as the halide, preferably the iodide derivative, of 2-propen-1-ol when R^{18} is 2-propenyl or 3-buten-1-ol when R^{18} is 3-butenyl, in the presence of a base, such as potassium carbonate, cesium carbonate or sodium hydride, preferably sodium hydride when R^{18} is 2-propenyl or cesium carbonate when R^{18} is 3-butenyl. The reaction is stirred in a polar solvent, such as dimethylformamide, at room temperature, for a time period between about 2 hours to about 48 hours, preferably about 18 hours.

In reaction 2 of Scheme 2, the compound of formula VIII is converted to the carboxylic acid compound of formula IV, wherein n is 2. The compound of formula VIII, wherein R^{18} is 2-propenyl, is converted to the compound of formula IV, wherein n is 2, by reacting VIII with borane-dimethylsulfide complex, followed by immediate oxidation using chromium trioxide in aqueous acetic acid. The oxidative

- 23 -

cleavage of terminal olefins to carboxylic acids can be achieved by several methods known in the art. The preferred method for the oxidative cleavage of the compound of formula VIII, wherein R^{18} is 3-butenyl, to obtain the carboxylic acid compound of formula IV is to
5 react VIII with sodium periodate in the presence of a catalytic amount of ruthenium (III) chloride in a mixture of carbon tetrachloride, acetonitrile and water.

10 The compound of formula IV, wherein n is 2, is further reacted to provide the hydroxamic acid compound of formula I, wherein n is 2, according to the procedure described above in reactions 4, 5 and 6 of Scheme 1.

15 An alternative method for the synthesis of the hydroxamic acid compound of formula I, wherein n is 1 and R^3 and R^4 are both hydrogen, is shown in reaction 1 of Scheme 3, beginning with reacting iminoacetic acid or a metal or ammonium salt of iminoacetic acid of
20 formula X with a functional derivative of an arylsulfonic acid compound, such as an arylsulfonyl chloride, at room temperature, in the presence of a suitable base, such as triethylamine, and a polar solvent such as tetrahydrofuran, dioxane, water or
25 acetonitrile, preferably a mixture of dioxane and water, to give the corresponding dicarboxylic acid compound of formula XI.

In reaction 2 of Scheme 3, the dicarboxylic acid compound of formula XI is dehydrated to give a
30 cyclic anhydride compound of formula XII. The formation of cyclic anhydrides by dehydration of dicarboxylic acids may be achieved by a variety of means. The preferred method for the dehydration of the dicarboxylic acid compound of formula XI to give a
35 cyclic anhydride compound of formula XII is to treat XI with an excess of acetic anhydride at a temperature between about 25°C. to about 80°C., preferably about

- 24 -

60°C. Excess acetic anhydride and acetic acid, a by-product of the reaction, are removed by evaporation under reduced pressure leaving the cyclic anhydride compound of formula XII.

5 In reaction 3 of Scheme 3, the cyclic anhydride compound of formula XII is reacted, at room temperature, with an amine, NR^1R^2 , or a salt of the amine, such as the hydrochloride, in the presence of a base, such as triethylamine, to give the carboxylic
10 acid of formula II, wherein n is 1 and R^3 and R^4 are both hydrogen. Suitable solvents for the reaction are those that will not react with the starting materials, which include chloroform, methylene chloride and dimethylformamide, preferably methylene chloride.


15 The compound of formula II is further reacted to give the hydroxamic acid compound of formula I, wherein n is 1 and R^3 and R^4 are both hydrogen, according to the procedure described above in reaction 6 of Scheme 1.

20 In reaction 1 of Scheme 4, the carboxylic acid compound of formula IV, wherein n is 2, is converted to the corresponding compound of formula V, wherein R^{19} is $(\text{C}_1\text{-C}_6)$ alkyl or tert-butyl, by reacting IV with a compound of the formula



wherein R^{19} is $(\text{C}_1\text{-C}_6)$ alkyl or tert-butyl, in an inert solvent, such as toluene, at a temperature between about 60°C. to about 100°C., preferably about 100°C., for a time period between about 1 hour to about 3
30 hours, preferably 2 hours. In reaction 2 of Scheme 4, the arylsulfonyl amino compound of formula VI wherein n is 1, 3, 4, 5 or 6 and R^{16} is $(\text{C}_1\text{-C}_6)$ alkyl, benzyl, allyl or tert-butyl, is converted to the corresponding compound of formula XIII, wherein R^{19} is $(\text{C}_1\text{-C}_6)$ alkyl
35 or tert-butyl, by reacting VI with a reactive derivativ of an alcohol of the formula

- 25 -



such as the chloride, bromide or iodide derivative, preferably the bromide derivative, wherein R^{19} is (C_1 - C_6)alkyl or tert-butyl, in the presence of a base such as potassium carbonate or sodium hydride, preferably sodium hydride, and a polar solvent, such as dimethylformamide. The reaction is stirred, at room temperature, for a time period between about 60 minutes to about 48 hours, preferably about 18 hours. The R^{16} protecting group, of the compounds of formulas IV and VI, is chosen such that it may be selectively removed in the presence of and without loss of the R^{19} (C_1 - C_6)alkyl or tert-butyl group, therefore, R^{16} cannot be the same as R^{19} . Removal of the R^{16} protecting group from the compound of formula XIII to give the corresponding carboxylic acid of formula XIV, wherein n is 1 to 6, in reaction 3 of Scheme 4, is carried out under conditions appropriate for that particular R^{16} protecting group in use which will not affect the R^{19} (C_1 - C_6)alkyl or tert-butyl group. Such conditions include; (a) saponification where R^{16} is (C_1 - C_6)alkyl and R^{19} is tert-butyl, (b) hydrogenolysis where R^{16} is benzyl and R^{19} is tert-butyl or (C_1 - C_6)alkyl, (c) treatment with a strong acid such as trifluoroacetic acid or hydrochloric acid where R^{16} is tert-butyl and R^{19} is (C_1 - C_6)alkyl, or (d) treatment with tributyltinhydride and acetic acid in the presence of catalytic bis(triphenylphosphine) palladium (II) chloride where R^{16} is allyl and R^{19} is (C_1 - C_6)alkyl or tert-butyl.

In reaction 4 of Scheme 4, the carboxylic acid of formula XIV is converted to the hydroxamic acid compound of formula XV, wherein n is 1 to 6, by treating XIV with 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide and 1-hydroxybenzotriazole in a polar solvent, such as dimethylformamide, followed by the

- 26 -

addition of hydroxylamine to the reaction mixture after a time period between about 15 minutes to about 1 hour, preferably about 30 minutes. The hydroxylamine is preferably generated in situ from a salt form, such as hydroxylamine hydrochloride, in the presence of a base, such as N-methylmorpholine. Alternatively, a protected derivative of hydroxylamine or its salt form, where the hydroxyl group is protected as a tert-butyl, benzyl or allyl ether, may be used in the presence of (benzotriazol-1-yloxy)tris(dimethylamino) phosphonium hexafluorophosphate and a base, such as N-methylmorpholine. Removal of the hydroxylamine protecting groups is carried out by hydrogenolysis for a benzyl protecting group or treatment with a strong acid, such as trifluoroacetic acid, for a tert-butyl protecting group. The allyl protecting group may be removed by treatment with tributyltinhydride and acetic acid in the presence of catalytic bis(triphenylphosphine) palladium (II) chloride. N,O-bis(4-methoxybenzyl)hydroxylamine may also be used, when R¹⁹ is (C₁-C₆)alkyl, as the protected hydroxylamine derivative where deprotection is achieved using a mixture of methanesulfonic acid and trifluoroacetic acid.

In reaction 5 of Scheme 4, the amide formula of formula XV is, if desired, converted to the corresponding carboxylic acid compound of formula XVI by (a) saponification where R¹⁹ is lower alkyl or (b) treatment with a strong acid, such as trifluoroacetic acid or hydrochloric acid, where R¹⁹ is tert-butyl.

Pharmaceutically acceptable salts of the acidic compounds of the invention are salts formed with bases, namely cationic salts such as alkali and alkaline earth metal salts, such as sodium, lithium, potassium, calcium, magnesium, as well as ammonium salts, such as ammonium, trimethyl-ammonium, diethylammonium, and tris-(hydroxymethyl)-

- 27 -

methylammonium salts.

Similarly acid addition salts, such as of mineral acids, organic carboxylic and organic sulfonic acids e.g. hydrochloric acid, methanesulfonic acid, maleic acid, are also possible provided a basic group, such as pyridyl, constitutes part of the structure.

The ability of the compounds of formula I or their pharmaceutically acceptable salts (hereinafter also referred to as the compounds of the present invention) to inhibit matrix metalloproteinases or the production of tumor necrosis factor (TNF) and, consequently, demonstrate their effectiveness for treating diseases characterized by matrix metalloproteinase or the production of tumor necrosis factor is shown by the following in vitro assay tests.

Biological Assay

Inhibition of Human Collagenase (MMP-1)

Human recombinant collagenase is activated with trypsin using the following ratio: 10 μ g trypsin per 100 μ g of collagenase. The trypsin and collagenase are incubated at room temperature for 10 minutes then a five fold excess (50 μ g/10 μ g trypsin) of soybean trypsin inhibitor is added.

10 mM stock solutions of inhibitors are made up in dimethyl sulfoxide and then diluted using the following Scheme:

10 mM ----> 120 μ M ----> 12 μ M ----> 1.2 μ M ----> 0.12 μ M

Twenty-five microliters of each concentration is then added in triplicate to appropriate wells of a 96 well microfluor plate. The final concentration of inhibitor will be a 1:4 dilution after addition of enzyme and substrate. Positive controls (enzyme, no inhibitor) are set up in wells D1-D6 and blanks (no enzyme, no inhibitors) are set in wells D7-D12.

Collagenase is diluted to 400 ng/ml and 25 μ l is then added to appropriate wells of the microfluor plate. Final concentration of collagenase in the assay

- 28 -

is 100 ng/ml.

Substrate (DNP-Pro-Cha-Gly-Cys(Me)-His-Ala-Lys(NMA)-NH₂) is made as a 5 mM stock in dimethyl sulfoxide and then diluted to 20 μM in assay buffer.

5 The assay is initiated by the addition of 50 μl substrate per well of the microfluor plate to give a final concentration of 10 μM.

Fluorescence readings (360 nm excitation, 460 nm emission) were taken at time 0 and then at 20 minute intervals. The assay is conducted at room temperature with a typical assay time of 3 hours.

Fluorescence vs time is then plotted for both the blank and collagenase containing samples (data from triplicate determinations is averaged). A time point that provides a good signal (the blank) and that is on a linear part of the curve (usually around 120 minutes) is chosen to determine IC₅₀ values. The zero time is used as a blank for each compound at each concentration and these values are subtracted from the 120 minute data. Data is plotted as inhibitor concentration vs % control (inhibitor fluorescence divided by fluorescence of collagenase alone x 100). IC₅₀'s are determined from the concentration of inhibitor that gives a signal that is 50% of the control.

25 If IC₅₀'s are reported to be <0.03 μM then the inhibitors are assayed at concentrations of 0.3 μM, 0.03 μM, 0.03 μM and 0.003 μM.

Inhibition of Gelatinase (MMP-2)

Inhibition of gelatinase activity is assayed using the Dnp-Pro-Cha-Gly-Cys(Me)-His-Ala-Lys(NMA)-NH₂ substrate (10 μM) under the same conditions as inhibition of human collagenase (MMP-1).

72kD gelatinase is activated with 1 mM APMA (p-aminophenyl mercuric acetate) for 15 hours at 4°C. and is diluted to give a final concentration in the assay of 100 mg/ml. Inhibitors are diluted as for inhibition of human collagenase (MMP-1) to give final

- 29 -

concentrations in the assay of 30 μM , 3 μM , 0.3 μM and 0.03 μM . Each concentration is done in triplicate.

Fluorescence readings (360 nm excitation, 460 emission) are taken at time zero and then at 20 minutes intervals for 4 hours.

IC₅₀'s are determined as per inhibition of human collagenase (MMP-1). If IC₅₀'s are reported to be less than 0.03 μM , then the inhibitors are assayed at final concentrations of 0.3 μM , 0.03 μM , 0.003 μM and 0.003 μM .

Inhibition of Stromelysin Activity (MMP-3)

Inhibition of stromelysin activity is based on a modified spectrophotometric assay described by Weingarten and Feder (Weingarten, H. and Feder, J., Spectrophotometric Assay for Vertebrate Collagenase, Anal. Biochem. 147, 437-440 (1985)). Hydrolysis of the thio peptolide substrate [Ac-Pro-Leu-Gly-SCH[CH₂CH(CH₃)₂]CO-Leu-Gly-OC₂H₅] yields a mercaptan fragment that can be monitored in the presence of Ellman's reagent.

Human recombinant prostromelysin is activated with trypsin using a ratio of 1 μl of a 10 mg/ml trypsin stock per 26 μg of stromelysin. The trypsin and stromelysin are incubated at 37°C. for 15 minutes followed by 10 μl of 10 mg/ml soybean trypsin inhibitor for 10 minutes at 37°C. to quench trypsin activity.

Assays are conducted in a total volume of 250 μl of assay buffer (200 mM sodium chloride, 50 mM MES, and 10 mM calcium chloride, pH 6.0) in 96-well microliter plates. Activated stromelysin is diluted in assay buffer to 25 $\mu\text{g}/\text{ml}$. Ellman's reagent (3-Carboxy-4-nitrophenyl disulfide) is made as a 1M stock in dimethyl formamide and diluted to 5 mM in assay buffer with 50 μl per well yielding at 1 mM final concentration.

10 mM stock solutions of inhibitors are made in dimethyl sulfoxide and diluted serially in assay

- 30 -

buffer such that addition of 50 μ L to the appropriate wells yields final concentrations of 3 μ M, 0.3 μ M, 0.003 μ M, and 0.0003 μ M. All conditions are completed in triplicate.

5 A 300 mM dimethyl sulfoxide stock solution of the peptide substrate is diluted to 15 mM in assay buffer and the assay is initiated by addition of 50 μ l to each well to give a final concentration of 3 mM substrate. Blanks consist of the peptide substrate and
10 Ellman's reagent without the enzyme. Product formation was monitored at 405 nm with a Molecular Devices UVmax plate reader.

IC₅₀ values were determined in the same manner as for collagenase.

15 Inhibition of MMP-13

Human recombinant MMP-13 is activated with 2mM APMA (p-aminophenyl mercuric acetate) for 1.5 hours, at 37°C. and is diluted to 400 mg/ml in assay buffer (50 mM Tris, pH 7.5, 200 mM sodium chloride, 5mM
20 calcium chloride, 20 μ M zinc chloride, 0.02% brij). Twenty-five microliters of diluted enzyme is added per well of a 96 well microfluor plate. The enzyme is then diluted in a 1:4 ratio in the assay by the addition of inhibitor and substrate to give a final concentration
25 in the assay of 100 mg/ml.

10 mM stock solutions of inhibitors are made up in dimethyl sulfoxide and then diluted in assay buffer as per the inhibitor dilution scheme for inhibition of human collagenase (MMP-1): Twenty-five
30 microliters of each concentration is added in triplicate to the microfluor plate. The final concentrations in the assay are 30 μ M, 3 μ M, 0.3 μ M and 0.03 μ M.

Substrate (Dnp-Pro-Gly-Cys(Me)-His-Ala-
35 Lys(NMA)-NH₂) is prepared as for inhibition of human collagenase (MMP-1) and 50 μ l is added to each well to give a final assay concentration of 10 μ M.

- 31 -

Fluorescence readings (360 nM excitation; 450 emission) are taken at time 0 and every 5 minutes for 1 hour.

Positive controls consist of enzyme and substrate with no inhibitor and blanks consist of substrate only.

IC₅₀'s are determined as per inhibition of human collagenase (MMP-1). If IC₅₀'s are reported to be less than 0.03 μ M, inhibitors are then assayed at final concentrations of 0.3 μ M, 0.03 μ M, 0.003 μ M and 0.0003 μ M.

Inhibition of TNF Production

The ability of the compounds or the pharmaceutically acceptable salts thereof to inhibit the production of TNF and, consequently, demonstrate their effectiveness for treating diseases involving the production of TNF is shown by the following in vitro assay:

Human mononuclear cells were isolated from anti-coagulated human blood using a one-step Ficoll-hypaque separation technique. The mononuclear cells were washed three times in Hanks balanced salt solution (HBSS) with divalent cations and resuspended to a density of 2×10^6 /ml in HBSS containing 1% BSA. Differential counts determined using the Abbott Cell Dyn 3500 analyzer indicated that monocytes ranged from 17 to 24% of the total cells in these preparations.

180 μ of the cell suspension was aliquoted into flat bottom 96 well plates (Costar). Additions of compounds and LPS (100ng/ml final concentration) gave a final volume of 200 μ l. All conditions were performed in triplicate. After a four hour incubation at 37°C. in an humidified CO₂ incubator, plates were removed and centrifuged (10 minutes at approximately 250 x g) and the supernatants removed and assayed for TNF α using the R&D ELISA Kit.

For administration to humans for the inhibition of matrix metalloproteinases or the

- 32 -

production of tumor necrosis factor (TNF), a variety of conventional routes may be used including orally, parenterally and topically. In general, the active compound will be administered orally or parenterally at dosages between about 0.1 and 25 mg/kg body weight of the subject to be treated per day, preferably from about 0.3 to 5 mg/kg. However, some variation in dosage will necessarily occur depending on the condition of the subject being treated. The person responsible for administration will, in any event, determine the appropriate dose for the individual subject.

The compounds of the present invention can be administered in a wide variety of different dosage forms, in general, the therapeutically effective compounds of this invention are present in such dosage forms at concentration levels ranging from about 5.0% to about 70% by weight.

For oral administration, tablets containing various excipients such as microcrystalline cellulose, sodium citrate, calcium carbonate, dicalcium phosphate and glycine may be employed along with various disintegrants such as starch (and preferably corn, potato or tapioca starch), alginic acid and certain complex silicates, together with granulation binders like polyvinylpyrrolidone, sucrose, gelatin and acacia. Additionally, lubricating agents such as magnesium stearate, sodium lauryl sulfate and talc are often very useful for tableting purposes. Solid compositions of a similar type may also be employed as fillers in gelatin capsules; preferred materials in this connection also include lactose or milk sugar as well as high molecular weight polyethylene glycols. When aqueous suspensions and/or elixirs are desired for oral administration, the active ingredient may be combined with various sweetening or flavoring agents, coloring matter or dyes, and, if so desired,

- 33 -

emulsifying and/or suspending agents as well, together with such diluents as water, ethanol, propylene glycol, glycerin and various like combinations thereof.

For parenteral administration (intramuscular, intraperitoneal, subcutaneous and intravenous use) a sterile injectable solution of the active ingredient is usually prepared. Solutions of a therapeutic compound of the present invention in either sesame or peanut oil or in aqueous propylene glycol may be employed. The aqueous solutions should be suitably adjusted and buffered, preferably at a pH of greater than 8, if necessary and the liquid diluent first rendered isotonic. These aqueous solutions are suitable for intravenous injection purposes. The oily solutions are suitable for intraarticular, intramuscular and subcutaneous injection purposes. The preparation of all these solutions under sterile conditions is readily accomplished by standard pharmaceutical techniques well known to those skilled in the art.

The present invention is illustrated by the following examples, but it is not limited to the details thereof.

EXAMPLE 1

2-(R)-N-Hydroxy-2-[(4-methoxybenzenesulfonyl)(2-morpholin-4-yl-2-oxoethyl)amino]-3-methylbutyramide

To a solution of D-valine benzyl ester hydrochloride (2.4 grams, 10 mmol) and triethylamine (2.5 grams, 3.5 mL, 25 mmol) in water (50 mL) and 1,4-dioxane (50 mL) is added 4-methoxybenzenesulfonyl chloride (2.3 grams, 11 mmol). The mixture was stirred at room temperature for 1 hour and then most of the solvent was removed by evaporation under vacuum. The mixture was diluted with ethyl acetate and was washed successively with dilute hydrochloric acid solution, water and brine. The organic solution was dried over magnesium sulfate and concentrated to leave N-(4-methoxybenzenesulfonyl)-D-valine benzyl ester as a

- 34 -

white solid, 3.6 grams (97%); m.p. 92-94°C.

N-(4-Methoxybenzenesulfonyl)-D-valine benzyl ester (1.50 grams, 4.0 mmol) was added to a suspension of sodium hydride (0.1 grams, 4.2 mmol) in dry dimethylformamide (20 mL) and, after 30 minutes, tert-butyl bromoacetate (0.8 mL, 4.2 mmol) was added. The resulting mixture was stirred overnight at room temperature and was then quenched by addition of saturated ammonium chloride solution (3 mL). The dimethylformamide was removed by evaporation under vacuum. The residue was taken up in ethyl acetate and washed with water and brine. After drying over magnesium sulfate, ethyl acetate was evaporated to leave an oil from which 2-(R)-2-[tert-butoxycarbonylmethyl(4-methoxybenzenesulfonyl)amino]-3-methylbutyric acid benzyl ester, a clear oil (1.92 grams, 98%), was isolated using flash chromatography on silica gel eluting with 15% ethyl acetate in hexane.

To a cold (0°C.) solution of 2-(R)-2-[tert-butoxycarbonylmethyl(4-methoxybenzenesulfonyl)amino]-3-methylbutyric acid benzyl ester (1.92 grams, 3.9 mmol) in methylene chloride (28 mL) was added trifluoroacetic acid (7 mL). The resulting solution was allowed to warm to room temperature and was stirred overnight. The methylene chloride and trifluoroacetic acid were evaporated under vacuum leaving 2-(R)-2-[carboxymethyl(4-methoxybenzenesulfonyl)amino]-3-methylbutyric acid benzyl ester as a clear oil, 1.70 grams (100%).

To a solution of 2-(R)-2-[carboxymethyl(4-methoxybenzenesulfonyl)amino]-3-methylbutyric acid benzyl ester (573 mg, 1.32 mmol) in methylene chloride (12 mL) were added sequentially triethylamine (0.46 mL, 3.28 mmol), morpholine (0.127 mL, 1.46 mmol) and (benzotriazol-1-yloxy)tris(dimethylamino)phosphonium hexafluorophosphate (646 mg, 1.46 mmol). The mixture was stirred at room temperature overnight and then

- 35 -

diluted with ethyl acetate. The solution was washed with 0.5 N hydrochloric acid solution and brine, dried over magnesium sulfate and concentrated under vacuum. The residue was chromatographed on silica gel using 40% ethyl acetate in hexane affording 2-(R)-2-[(4-methoxybenzenesulfonyl)(2-morpholin-4-yl-oxoethyl)amino]-3-methylbutyric acid benzyl ester as a clear oil, 590 mg (89%).

To a solution of 2-(R)-2-[(4-methoxybenzenesulfonyl)(2-morpholin-4-yl-2-oxoethyl)amino]-3-methylbutyric acid benzyl ester (590 mg, 1.17 mmol) in ethanol (50 mL) was added 10% palladium on activated carbon (200 mg). The mixture was agitated under 3 atmospheres hydrogen in a Parr shaker for 2 hours. The catalyst was removed by filtration through nylon (pore size 0.45 μ m) and the solvent was evaporated leaving 2-(R)-2-[(4-methoxybenzenesulfonyl)(2-morpholin-4-yl-2-oxoethyl)amino]-3-methylbutyric acid as a white foam, 485 mg (100%).

To a solution of 2-(R)-2-[(4-methoxybenzenesulfonyl)(2-morpholin-4-yl-2-oxoethyl)amino]-3-methylbutyric acid (485 mg, 1.17 mmol) in methylene chloride (12 mL) were added sequentially triethylamine (0.52 mL, 3.71 mmol), O-benzylhydroxylamine hydrochloride (205 mg, 1.28 mmol) and (benzotriazol-1-yloxy)tris(dimethylamino)phosphonium hexafluorophosphate (570 mg, 1.29 mmol). The mixture was stirred at room temperature overnight and then diluted with ethyl acetate. The solution was washed sequentially with 0.5 N hydrochloric acid solution, water, saturated sodium hydrogen carbonate solution and brine, dried over magnesium sulfate and concentrated under vacuum. The residue was chromatographed on silica gel using 20% hexane in ethyl acetate to afford 2-(R)-N-benzyloxy-2-[(4-methoxybenzenesulfonyl)(2-

- 36 -

morpholin-4-yl-2-oxoethyl)amino]-3-methylbutyramide as a white foam, 510 mg (84%).

To a solution of 2-(R)-N-benzyloxy-2-[(4-methoxybenzenesulfonyl)(2-morpholin-4-yl-2-oxoethyl)amino]-3-methylbutyramide (510 mg, 0.98 mmol) in methanol (50 mL) was added 5% palladium on activated carbon (120 mg). The mixture was agitated under 2 atmospheres hydrogen in a Parr shaker for 2 hours. The catalyst was removed by filtration through nylon (pore size 0.45 μ m) and the solvent was evaporated leaving 2-(R)-N-hydroxy-2-[(4-methoxybenzenesulfonyl)(2-morpholin-4-yl-2-oxoethyl)amino]-3-methylbutyramide as a white solid, 418 mg (99%); ^1H NMR (CDCl_3): δ 10.3 (br s, 1H), 7.90 (br s, 1H, overlapped), 7.86 (d, J = 8.8 Hz, 2H, overlapped), 6.94 (d, J = 8.8 Hz, 2H), 4.39 (d, J = 17.1 Hz, 1H), 4.09 (d, J = 17.1, 1H), 3.84 (s, 3H), 3.80-3.48 (m, 9H), 2.20-1.95 (m, 1H), 0.82 (d, J = 6.5 Hz, 3H), 0.45 (d, J = 6.5 Hz, 3H); MS (LSIMS): m/z 430 (M+H).

20

EXAMPLE 2

2-(R)-N-Hydroxy-2-[(4-methoxybenzenesulfonyl)(3-morpholin-4-yl-3-oxopropyl)amino]-3-methylbutyramide

To a solution of N-(4-methoxybenzenesulfonyl)-D-valine benzyl ester (2.2 grams, 5.83 mmol) in dry dimethylformamide (40 mL) were added cesium carbonate (2.3 grams, 7.1 mmol) and 1-iodo-3-butene (1.3 grams, 7.1 mmol). The mixture was stirred at room temperature overnight and was then poured into water. The mixture was extracted twice with ether and the combined ether extracts were washed with brine, dried over magnesium sulfate and concentrated under reduced pressure. The residue was taken up in 20% ethyl acetate/hexane; starting material N-(4-methoxybenzenesulfonyl)-D-valine benzyl ester (1.5 g) crystallized from the mixture and was recovered by filtration. The filtrate was concentrated under vacuum and the residue was chromatographed on silica

- 37 -

gel using 20% ethyl acetate/hexane as eluent to provide 2-(R)-2-[but-3-enyl(4-methoxybenzenesulfonyl)amino]-3-methylbutyric acid benzyl ester as a clear oil, 404 mg (16%).

5 To a mixture of 2-(R)-2-[but-3-enyl(4-methoxybenzenesulfonyl)amino]-3-methylbutyric acid benzyl ester (780 mg, 1.81 mmol) and ruthenium (III) chloride hydrate (10 mg, 0.048 mmol) in acetonitrile (6 mL), carbon tetrachloride (6 mL) and water (8 mL) was
10 added sodium periodate (1.7 grams, 7.9 mmol). After stirring at room temperature for 2 hours, the mixture was diluted with methylene chloride and filtered through diatomaceous earth. The organic layer was
15 separated, washed with dilute hydrochloric acid solution and brine, dried over magnesium sulfate and concentrated to leave 2-(R)-2-[2-carboxyethyl(4-methoxybenzenesulfonyl)amino]-3-methylbutyric acid benzyl ester as an oil, 710 mg (87%).

Alternatively, the intermediate 2-(R)-2-[2-carboxyethyl(4-methoxybenzenesulfonyl)amino]-3-methylbutyric acid benzyl ester was prepared by the
20 following higher yielding procedure:

N-(4-Methoxybenzenesulfonyl)-D-valine benzyl ester (18.8 grams, 49.8 mmol) was added to a suspension
25 of sodium hydride (1.3 grams, 54 mmol) in dry dimethylformamide (200 mL) and, after 1.5 hours, a solution of allyl bromide (4.7 mL, 54 mmol) was added. The resulting mixture was stirred overnight at room temperature and was then quenched by addition of
30 saturated ammonium chloride solution. The dimethylformamide was removed by evaporation under vacuum. The residue was taken up in ether and washed with water and brine. After drying over magnesium sulfate, ether was evaporated to leave an oil from
35 which 2-(R)-2-[(4-methoxybenzenesulfonyl)prop-2-enylamino]-3-methylbutyric acid benzyl ester, a clear oil (18.1 grams, 87%), was isolated using flash

- 38 -

chromatography on silica gel eluting with 10% ethyl acetate in hexane and then 20% ethyl acetate in hexane.

To a 1 M solution of borane/disulfide complex in methylene chloride (1.45 mL, 2.9 mmol) was added a solution of 2-(R)-2-[(4-methoxybenzenesulfonyl)prop-2-enylamino]-3-methylbutyric acid benzyl ester (3.6 grams, 8.6 mmol) in methylene chloride (8 mL). The solution was stirred at room temperature for 4 hours at which time more 1 M solution of borane/disulfide complex in methylene chloride (2.0 mL, 4.0 mmol) was added. The mixture was stirred at room temperature overnight and was then added dropwise to a mechanically stirred solution of chromium trioxide (5.1 grams, 51.6 mole) in acetic acid (31 mL) and water (3.5 mL) while keeping the internal temperature between -5°C. and 10°C. After stirring at room temperature overnight, the mixture was diluted with water and extracted with methylene chloride. The extract was washed with brine, dried (magnesium sulfate) and concentrated. The residue was chromatographed on silica gel eluting successively with chloroform and 2% methanol in chloroform to afford 2-(R)-2-[2-carboxyethyl(4-methoxybenzenesulfonyl)amino]-3-methylbutyric acid benzyl as an oil (2.42 grams, 63%).

To a solution of 2-(R)-2-[2-carboxyethyl(4-methoxybenzenesulfonyl)amino]-3-methylbutyric acid benzyl ester (710 mg, 1.58 mmol) in methylene chloride (15 mL) were added sequentially triethylamine (0.47 mL, 3.35 mmol), morpholine (0.15 mL, 1.72 mmol) and (benzotriazol-1-yloxy)tris(dimethylamino)phosphonium hexafluorophosphate (769 mg, 1.74 mmol). The mixture was stirred at room temperature overnight and then diluted with methylene chloride. The solution was washed with 0.5 N hydrochloric acid solution and brine, dried over magnesium sulfate and concentrated under vacuum. The solid residue was chromatographed on silica gel using 20% hexane in ethyl acetate affording

- 39 -

2-(R)-2-[(4-methoxybenzenesulfonyl)(3-morpholin-4-yl-3-oxopropyl)amino]-3-methylbutyric acid benzyl ester as a clear oil, 725 mg (88%).

To a solution of 2-(R)-2-[(4-methoxybenzenesulfonyl)(3-morpholin-4-yl-3-oxopropyl)amino]-3-methylbutyric acid benzyl ester (725 mg, 1.40 mmol) in ethanol (35 mL) was added 10% palladium on activated carbon (50 mg). The mixture was agitated under 3 atmospheres hydrogen in a Parr shaker for 3 hours. The catalyst was removed by filtration through nylon (pore size 0.45 μ m) and the solvent was evaporated leaving 2-(R)-(2)-[(4-methoxybenzenesulfonyl)(3-morpholin-4-yl-3-oxopropyl)amino]-3-methyl-butyrlic acid as a white solid, 540 mg (90%).

To a solution of 2-(R)-2-[(4-methoxybenzenesulfonyl)(3-morpholin-4-yl-3-oxopropyl)amino]-3-methylbutyric acid (540 mg, 1.26 mmol) and 1-hydroxybenztriazole hydrate (205 mg, 1.33 mmol) in dry dimethylformamide (12 mL) was added 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride (289 mg, 1.51 mmol). After stirring for 30 minutes, hydroxylamine hydrochloride (350 mg, 5.04 mmol) and then triethylamine (1.0 mL, 7.17 mmol) were added. The mixture was stirred at room temperature overnight and then diluted with ethyl acetate. The solution was washed sequentially with water, 0.5 N hydrochloric acid solution and brine. The solution was then dried over magnesium sulfate and concentrated under vacuum to leave a white foam. The material was dissolved in toluene, filtered and concentrated. The residue was triturated with ether to afford 2-(R)-N-hydroxy-2[(4-methoxybenzenesulfonyl)(3-morpholin-4-yl-3-oxopropyl)amino]-3-methylbutyramide as a solid, 200 mg (36%); ^1H NMR (CDCl_3): δ 9.35 (br s, 1H), 7.73 (d, J = 8.9 Hz, 2H), 6.95 (d, J = 8.9 Hz, 2H), 3.86 (s, 3H), 3.83-3.73 (m, 1H), 3.70-3.52 (m, 7H), 3.46-3.43 (m,

- 40 -

2H), 3.41-3.29 (m, 1H), 2.92-2.69 (m, 2H), 2.30-2.17 (m, 1H), 0.84 (d, J = 6.5 Hz, 3H), 0.41 (d, J = 6.5 Hz, 3H); MS (particle beam): m/z 444 (M+H), 428, 383, 329; HRMS calculated for C₁₉H₃₀N₃O₇S (M+H): 444.1804,

5 Found: 464.1818.

The title compounds of Examples 3-6 were prepared by a method analogous to that described in Example 2 using 2-(R)-2-[2-carboxyethyl(4-methoxybenzenesulfonyl)amino]-3-methylbutyric acid
10 benzyl ester as the starting material which is coupled with the amine indicated.

EXAMPLE 3

2-(R)-2-[(2-Benzylcarbamoyl)ethyl](4-methoxybenzenesulfonyl)amino]-N-hydroxy-3-methylbutyramide
15

Coupled with benzylamine; ¹H NMR (DMSO-d₆):
δ 10.72 (s, 1H), 8.89 (s, 1H), 8.39 (m, 1H), 7.74 (d, J = 9.0 Hz, 2H), 7.32-7.21 (m, 5H), 7.05 (d, J = 9.0 Hz, 2H), 4.21 (d, J = 5.9 Hz, 2H), 4.02-3.87 (m, 1H), 3.82
20 (s, 3H), 3.63 (d, J = 10.8 Hz, 1H), 3.29-3.17 (m, 1H), 2.71-2.57 (m, 1H), 2.52-2.40 (m, 1H), 2.06-1.94 (m, 1H), 0.77 (d, J = 6.6 Hz, 3H), 0.74 (d, J = 6.5 Hz, 3H); MS (LSIMS): m/z 464 (M+H); HRMS calculated for C₂₂H₃₀N₃O₆S (M+H): 464.1855. Found: 464.1832.

25

EXAMPLE 4

2-(R)-N-Hydroxy-2-[(4-methoxybenzenesulfonyl)(2-[(pyridin-3-ylmethyl)carbamoyl]ethyl)amino]-3-methylbutyramide

Coupled with 3-pyridylmethylamine: ¹H NMR (DMSO-d₆):
30 δ 10.72 (s, 1H), 8.89 (s, 1H), 8.49-8.42 (m, 3H), 7.73 (d, J = 8.9 Hz, 2H), 7.63-7.60 (m, 1H), 7.32 (dd, J = 4.8, 7.8 Hz, 1H), 7.05 (d, J = 8.9 Hz, 2H), 4.23 (d, J = 5.8 Hz, 2H), 4.00-3.88 (m, 1H), 3.81 (s, 3H), 3.62 (d, J = 10.8 Hz, 1H), 3.27-3.17 (m, 1H),
35 2.69-2.58 (m, 1H), 2.52-2.41 (m, 1H), 2.07-1.94 (m, 1H), 0.76 (d, J = 6.5 Hz, 3H), 0.72 (d, J = 6.4 Hz, 3H); MS (LSIMS): m/z 465 (M+H).

- 41 -

EXAMPLE 5

2-(R)-N-Hydroxy-2-([4-methoxybenzenesulfonyl]
[2-(methylpyridin-3-ylmethylcarbamoyl)ethyl]amino)-3-
methylbutyramide

5 Coupled with 3-(N-methylaminomethyl)pyridine:
¹H NMR (DMSO-d₆): δ 10.75 (br s, 1H), 8.92 (s, 1H),
8.52-8.29 (m, 2H), 7.75 (d, J = 8.8 Hz, 1.4 H), 7.67
(d, J = 8.8 Hz, 0.6 H), 7.62-7.58 (m, 1H), 7.42-7.32
(m, 1H), 7.06 (d, J = 8.8 Hz, 1.4 H), 7.01 (d, J = 8.8
10 Hz, 0.6 H), 4.55-4.41 (m, 2H), 3.94-3.82 (m, 1H), 3.81
(s, 2.1 H), 3.80 (s, 0.9 H), 3.68-3.60 (m, 1H), 3.33-
3.19 (m, 1H), 2.90-2.50 (m, 2H), 2.88 (s, 2.1 H
overlapped), 2.79 (s, 0.9 H), 2.05-1.80 (m, 1H),
0.79-0.63 (m, 6H): MS (thermospray): m/z 479
15 (M+H), 364.

EXAMPLE 6

4-(3-[(1-(R)-1-Hydroxycarbamoyl-2-
methylpropyl)(4-methoxybenzenesulfonyl)amino]propionyl)
piperazine-1-carboxylic acid, tert-butyl ester

20 Coupled with tert-butyl-1-
piperazinecarboxylate: ¹H NMR (DMSO-d₆): δ 10.77 (br
s, 1H), 8.93 (s, 1H), 7.74 (d, J = 8.9 Hz, 2H), 7.06
(d, J = 8.9 Hz, 2H), 3.90-3.80 (m, 1H), 3.82 (s, 3H,
overlapped), 3.64 (d, J = 10.8 Hz, 1H), 3.60-3.16 (m,
25 9H), 2.80-2.71 (m, 1H), 2.59-2.47 (m, 1H), 2.03-1.91
(m, 1H), 1.39 (s, 9H), 0.77 (d, J = 6.5 Hz, 3H), 0.71
(d, J = 6.5, 3H); MS (thermospray): m/z 543 (M+H),
443, 382, 328.

EXAMPLE 7

30 2-(R)-N-Hydroxy-2-[(4-methoxybenzenesulfonyl]
(3-oxo-3-piperazin-1-ylpropyl)amino]-3-methylbutyramide
hydrochloride

A solution of 4-(3-[(1-(R)-1-
hydroxycarbamoyl-2-methylpropyl)(4-
35 methoxybenzenesulfonyl)amino]propionyl)piperazine-1-
carboxylic acid, tert-butylester [Example 6] (430 mg,
0.79 mmol) in methylene chloride (11 mL) was cooled to

- 42 -

0°C. Hydrogen chloride gas was then bubbled through the solution for about 0.5 minute. The solution was allowed to warm to room temperature with stirring over 1 hour. Volatiles were removed by evaporation and the residue was filtered washing with methylene chloride to collect solid 2-(R)-N-hydroxy-2-[(4-methoxybenzenesulfonyl)(3-oxo-3-piperazin-1-ylpropyl)amino]-3-methylbutyramide hydrochloride, 375 mg (99%). ¹H NMR (DMSO-d₆): δ 10.78 (br s, 1H), 9.16 (br s, 1H), 7.74 (d, J = 8.8 Hz, 2H), 7.07 (d, J = 8.9 Hz, 2H), 3.82 (s, 3H), 3.62 (br s, 4H), 3.38-3.18 (m, 1H), 3.16-3.07 (br s, 2H), 3.07-2.98 (br s, 2H), 2.83-2.73 (m, 1H), 2.65-2.53 (m, 1H), 2.06-1.90 (m, 1H), 0.76 (d, J = 6.5 Hz, 3H), 0.72 (d, J = 6.5 Hz, 3H). A broad water peak between δ 4.0 and 3.5 obscured some signals from this compound; MS (thermospray): m/z 443 (M+H), 382, 328.

EXAMPLE 8

2-(R)-2-[(Benzylcarbamoylmethyl)(4-methoxybenzenesulfonyl)amino]-N-hydroxy-3-methylbutyramide

To a solution of 2-(R)-2-[carboxymethyl(4-methoxybenzenesulfonyl)amino]-3-methylbutyric acid benzyl ester (example 1) (905 mg, 2.08 mmol) in methylene chloride (18 mL) were added sequentially triethylamine (0.72 mL, 5.14 mmol), benzylamine (0.25 mL, 2.29 mmol) and (benzotriazol-1-yloxy)tris(dimethylamino)phosphonium hexafluorophosphate (1.01 grams, 2.28 mmol). The mixture was stirred at room temperature overnight and then diluted with ethyl acetate. The solution was washed with 0.5 N hydrochloric acid solution and brine, dried over magnesium sulfate and concentrated under vacuum. The residue was chromatographed on silica gel using a 2:5:16 ratio of methylene chloride/ethyl acetate/hexane affording 2-(R)-2-[(benzylcarbamoylmethyl)(4-methoxybenzenesulfonyl)amino]-3-methylbutyric acid

- 43 -

benzyl ester as a clear oil, 933 mg (86%).

To a solution of 2-(R)-2-

5 [(benzylcarbamoylmethyl)(4-methoxybenzenesulfonyl)-amino]-3-methylbutyric acid benzyl ester (933 mg, 1.17 mmol) in ethanol (50 mL) was added 10% palladium on activated carbon (85 mg). The mixture was agitated under 3 atmospheres hydrogen in a Parr shaker for 4 hours. The catalyst was removed by filtration through nylon (pore size 0.45 μ m) and the solvent was
10 evaporated leaving 2-(R)-2-[(benzylcarbamoylmethyl)(4-methoxybenzenesulfonyl)amino]-3-methylbutyric acid as a white foam, 755 mg (98%).

To a solution of 2-(R)-2-

15 [(benzylcarbamoylmethyl)(4-methoxybenzenesulfonyl)amino]-3-methylbutyric acid (655 mg, 1.51 mmol) and 1-hydroxybenztriazole hydrate (224 mg, 1.46 mmol) in dry dimethylformamide (15 mL) was added 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride (316 mg, 1.65 mmol). After stirring for
20 30 minutes, hydroxylamine hydrochloride (416 mg, 6.0 mmol) and then N-methylmorpholine (0.99 mL, 9.0 mmol) were added. The mixture was stirred at room temperature overnight and then diluted with ethyl acetate. The solution was washed sequentially with
25 water, 0.5 N hydrochloric acid solution and brine. The solution was then dried over magnesium sulfate and concentrated under vacuum to leave a white foam which was chromatographed on silica gel eluting with ethyl acetate to afford 2-(R)-2-[(benzylcarbamoylmethyl)(4-
30 methoxybenzene-sulfonyl)amino]-N-hydroxy-3-methylbutyramide as a white foam, 570 mg (84%); ^1H NMR (DMSO- d_6): δ 10.75 (br s, 1H), 8.90 (s, 1H), 8.47 (m, 1H), 7.85 (d, J = 8.9 Hz, 2H), 7.83-7.19 (m, 5H), 7.04 (d, J = 8.9 Hz, 2H), 4.37 (d, J = 11.4 Hz, 1H), 4.28
35 (d, J = 5.9 Hz, 2H), 3.89 (d, J = 11.4 Hz, 1H), 3.82 (s, 3H), 3.45 (d, J = 10.3 Hz, 1H), 1.90-1.79 (m, 1H), 0.73 (d, J = 6.3 Hz, 6H); MS (LSIMS): m/z 450 (M+H).

- 44 -

EXAMPLE 92-(R)-2-[(Benzylmethylcarbamoylmethyl)(4-methoxybenzenesulfonyl)amino]-N-hydroxy-3-methylbutyramide

5 To a solution of 2-(R)-2-[carboxymethyl(4-methoxybenzenesulfonyl)amino]-3-methylbutyric acid benzyl ester (Example 1) (1.05 grams, 2.41 mmol) in methylene chloride (20 mL) were added sequentially triethylamine (0.84 mL, 6.0 mmol), N-benzylmethylanine
10 (0.34 mL, 2.63 mmol) and (benzotriazol-1-ylloxy)tris(dimethylamino)phosphonium hexafluorophosphate (1.17 grams, 2.69 mmol). The mixture was stirred at room temperature overnight and then diluted with ethyl acetate. The solution was
15 washed with 0.5 N hydrochloric acid solution and brine, dried over magnesium sulfate and concentrated under vacuum. The residue was chromatographed on silica gel using 35% ethyl acetate in hexane (plus a small amount of methylene chloride to load the sample on the column)
20 affording 2-(R)-2-[benzylmethylcarbamoylmethyl)(4-methoxybenzenesulfonyl)amino]-3-methylbutyric acid benzyl ester as a clear oil, 1.14 grams (88%).

To a solution of 2-(R)-2-[(benzylmethylcarbamoylmethyl)(4-methoxybenzenesulfonyl)amino]-3-methylbutyric acid
25 benzyl ester (1.14 grams, 2.12 mmol) in ethanol (100 mL) was added 10% palladium on activated carbon (400 mg). The mixture was agitated under 3 atmospheres hydrogen in a Parr shaker for 3 hours. The catalyst
30 was removed by filtration through nylon (pore size 0.45 µm) and the solvent was evaporated leaving 2-(R)-2-[(benzylmethylcarbamoylmethyl)(4-methoxybenzenesulfonyl)amino]-3-methylbutyric acid as a white foam, 902 mg (95%).

35 To a solution of 2-(R)-2-[(benzylmethylcarbamoylmethyl)(4-methoxybenzenesulfonyl)amino]-3-methylbutyric acid (902

- 45 -

mg, 2.01 mmol) in methylene chloride (20 mL) were added sequentially triethylamine (0.90 mL, 6.42 mmol), O-allylhydroxylamine hydrochloride (242 mg, 2.21 mmol) and (benzotriazol-1-yl)oxytris-

5 (dimethylamino)phosphonium hexafluorophosphate (978 mg, 2.21 mmol). The mixture was stirred at room temperature overnight and then diluted with ethyl acetate. The solution was washed with 0.5 N hydrochloric acid solution and brine, dried over

10 magnesium sulfate and concentrated under vacuum. The residue was chromatographed on silica gel using 40% hexane in ethyl acetate to afford 2-(R)-N-allyloxy-2-[(benzylmethylcarbamoylmethyl)(4-methoxybenzenesulfonyl)amino]-3-methylbutyramide as an oil,

15 1.008 grams (100%).

To a solution of 2-(R)-N-allyloxy-2-[(benzylmethylcarbamoylmethyl)(4-methoxybenzenesulfonyl)amino]-3-methylbutyramide (500 mg, 0.99 mmol) in methylene chloride (40 mL) was added

20 bis(triphenylphosphine)palladium (II) chloride (280 mg, 0.4 mmol) and then, dropwise, tributyltinhydride (0.43 mL, 2.2 mmol). The solution was stirred at room temperature for 1 hour, diluted with methylene chloride, washed with 1 N hydrochloric acid solution,

25 dried over magnesium sulfate and concentrated. The residue was taken up in ethyl acetate and filtered to remove a solid. After concentration, the filtrate was chromatographed on silica gel eluting with chloroform and then 2% methanol in chloroform to afford 2-(R)-2-[(benzylmethylcarbamoylmethyl)(4-methoxybenzenesulfonyl)amino]-N-hydroxy-3-methylbutyramide as a white

30 solid (340 mg, 74%). ¹H NMR (DMSO-d₆): δ 10.66 (br s, 1H), 8.87 (br s, 0.6 H), 8.84 (s, 0.4 H), 7.91 (d, J = 8.9 Hz, 1.2 H), 7.78 (d, J = 8.9 Hz, 0.8 H), 7.43-7.21 (m, 5H), 7.05 (d, J = 8.9 Hz, 1.2 H), 7.00 (d, J = 8.9 Hz, 0.8 H), 4.72 (d, J = 17.7 Hz, 0.4 H), 4.70 (d, J = 17.7 Hz, 0.6 H), 4.59-4.42 (m, 1H), 4.25 (d, J = 17.8

35

- 46 -

Hz, 0.6H), 4.07 (d, J = 17.7 Hz, 0.4H), 3.82 (s, 3H),
3.46-3.40 (m, 1H), 2.91 (s, 1.8H), 2.83 (s, 1.2 H),
1.92-1.70 (m, 1H), 0.75-0.69 (m, 6H); MS (thermospray):
m/z 464 (M+H), 307, 239.

5 The title compounds of Examples 10-11 were
prepared by a method analogous to that described in
Example 9 using 2-(R)-2-[carboxymethyl(4-
methoxybenzenesulfonyl)amino]-3-methylbutyric acid
benzyl ester (Example 1) as the starting material which
10 is coupled with the amine indicated.

EXAMPLE 10

2-(R)-N-Hydroxy-2-[(4-
methoxybenzenesulfonyl)-[(2-morpholin-4-
ylethylcarbamoyl)methyl]amino]-3-methylbutyramide

15 Coupled with 4-(2-aminoethyl)morpholine: ¹H
NMR (DMSO-d₆): δ 10.74 (br s, 1H), 8.90 (br s, 1H),
7.84 (br s, 1H, overlapped), 7.84 (d, J = 8.8 Hz, 2H),
7.06 (d, J = 8.8 Hz, 2H), 4.33 (d, J = 17.5 Hz, 1H),
3.83 (s, 3H), 3.78 (d, J = 17.5 Hz, 1H), 3.57-3.54
20 (m, 4H), 3.49 (d, J = 10.2 Hz, 1H), 3.28-3.06 (m, 2H),
2.34-2.30 (m, 6H), 1.93-1.77 (m, 1H), 0.77-0.74
(m, 6H).

EXAMPLE 11

2-(R)-N-Hydroxy-2-[(4-
methoxybenzenesulfonyl)(2-oxo-2-pyrrolidin-1-
ylethyl)amino]-3-methylbutyramide

25 Coupled with pyrrolidine: ¹H NMR (CD₃OD): δ
7.90 (d, J = 8.9 Hz, 2H), 7.04 (d, J = 8.9 Hz, 2H),
4.50 (d, J = 17.6 Hz, 1H), 4.15 (d, J = 17.6 Hz, 1H),
30 3.87 (s, 3H), 3.56-3.39 (m, 5H), 2.07-1.82 (m, 5H),
0.83 (d, J = 6.6 Hz, 3H), 0.73 (d, J = 6.6 Hz, 3H); MS
(thermospray): m/z 414 (M+1); HRMS calculated for
C₁₈H₂₈N₃O₆S (M+H): 414.1699. Found 414.1703.

- 47 -

EXAMPLE 122-[Dimethylcarbamoylmethyl(4-methoxybenzenesulfonyl)amino]-N-hydroxy-3-methylbutyramide

5 A solution of 2-(R)-2-[carboxymethyl(4-methoxybenzenesulfonyl)amino]-3-methylbutyric acid benzyl ester (Example 1) (1.89 grams, 4.34 mmol) in thionyl chloride (25 mL) was heated at reflux for 1 hour. After cooling, the excess thionyl chloride was
10 evaporated. The residue was taken up in methylene chloride (50 mL) and the solution was cooled in an ice bath. Dimethylamine gas was slowly bubbled through the solution for 1 hour. After evaporation of the solvent, the residue was taken up in ethyl acetate, washed with
15 1 N hydrochloric acid solution, water and brine, dried over magnesium sulfate and concentrated to leave dimethylcarbamoylmethyl(4-methoxybenzenesulfonyl)amino-3-methylbutyric acid benzyl ester as an oil, 1.77 grams (88%).

20 To a solution of dimethylcarbamoylmethyl(4-methoxybenzenesulfonyl)amino-3-methylbutyric acid benzyl ester (1.77 grams, 3.83 mmol) in ethanol (100 mL) was added 10% palladium on activated carbon (644 mg). The mixture was agitated under 3 atmospheres
25 hydrogen in a Parr shaker for 1.5 hours. The catalyst was removed by filtration through nylon (pore size 0.45 μ m) and the solvent was evaporated leaving dimethylcarbamoylmethyl(4-methoxybenzenesulfonyl)amino-3-methylbutyric acid as a white foam, 1.42 grams (100%).

30 To a solution of dimethylcarbamoylmethyl(4-methoxybenzenesulfonyl)amino-3-methylbutyric acid (1.42 grams, 3.81 mmol) and 1-hydroxybenzotriazole hydrate (687 mg, 4.48 mmol) in dry dimethylformamide (7 mL) was added 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide
35 hydrochloride (974 mg, 5.08 mmol). After stirring for 30 minutes, hydroxylamine hydrochloride (1.17 grams, 16.8 mmol) and then N-methylmorpholine (2.8 mL, 25.5

- 48 -

mmol) were added. The mixture was stirred at room temperature overnight and then concentrated under vacuum. The residue was taken up in ethyl acetate and the resulting solution was washed sequentially with water, 0.5 N hydrochloric acid solution and brine. The solution was then dried over magnesium sulfate and concentrated under vacuum to leave an oil which was chromatographed on silica gel eluting successively with ethyl acetate, 5% methanol in chloroform and 10% methanol in chloroform to afford 2-[dimethyl-carbamoylmethyl(4-methoxybenzenesulfonyl)amino]-N-hydroxy-3-methylbutyramide as a white solid, 390 mg (26%). ¹H NMR (DMSO-d₆): δ 10.70 (br s, 1H), 8.89 (s, 1H), 7.80 (d, J = 8.9 Hz, 2H), 7.10 (d, J = 8.9 Hz, 2H), 4.62 (d, J = 17.7 Hz, 1H), 4.14 (d, J = 17.7 Hz, 1H), 3.84 (s, 3H), 3.40 (d, J = 10.4 Hz, 1H), 2.97 (s, 3H), 2.82 (s, 3H), 1.88-1.72 (m, 1H), 0.72 (d, J = 6.5 Hz, 6H); MS (thermospray): m/z 388 (M+1); HRMS calculated for C₁₆H₂₆N₃O₆S (M+H): 388.1542 Found: 388.1592.

EXAMPLE 13

2-(R)-2-N-Hydroxy-2-((4-methoxybenzenesulfonyl)((pyridin-3-ylmethyl)carbamoyl)methyl)amino)-3-methylbutyramide

2-(R)-N-Hydroxy-2-((4-methoxybenzenesulfonyl)((pyridin-3-ylmethyl)carbamoyl)methyl)amino)-3-methylbutyramide was prepared by a procedure similar to that of Example 12 starting with 2-(R)-2-[carboxymethyl(4-methoxybenzenesulfonyl)amino]-3-methylbutyric acid benzyl ester (Example 1) and coupling this to 3-pyridylmethylamine via the acid chloride. ¹H NMR (CD₃OD): δ 8.55-8.53 (m, 1H), 8.43-8.40 (m, 1H), 7.90-7.82 (m, 1H, overlapped), 7.86 (d, J = 8.9 Hz, 2H), 7.40 (dd, J = 4.8, 7.8 Hz, 1H), 7.04 (d, J = 8.9 Hz, 2H), 4.50 (d, J = 17.5 Hz, 1H), 4.39 (d, J = 17.5 Hz, 1H), 4.32 (d, J = 17.7 Hz, 1H), 4.02 (d, J = 17.7 Hz,

- 49 -

1H), 3.87 (s, 3H), 3.60 (d, J = 10.3 Hz, 1H), 2.08-1.93 (m, 1H), 0.85 (d, J = 6.5 Hz, 3H), 0.70 (d, J = 6.5 Hz, 3H); MS (thermospray): m/z 451 (M+H), 336, 320.

EXAMPLE 14

5 N-Hydroxy-[(4-methoxybenzenesulfonyl)(2-morpholin-4-yl-2-oxoethyl)amino]acetamide

To a solution of iminoacetic acid disodium salt monohydrate (5.0 grams, 25.6 mmol) in dioxane (50 ml) and water (50 ml) was added triethylamine (5.3 ml, 38 mmol) followed by 4-methoxybenzenesulfonyl chloride (5.8 grams, 28.0 mmol). The mixture was stirred overnight at room temperature and diluted with methylene chloride. The solution was washed with 1 N hydrochloric acid solution, water and brine, dried over magnesium sulfate and concentrated under vacuum leaving [carboxymethyl(4-methoxybenzenesulfonyl)amino]acetic acid as a white solid, 3.83 grams (49%).

[Carboxymethyl(4-methoxybenzenesulfonyl)amino]acetic acid (0.5 grams, 1.65 mmol) in acetic anhydride (15 mL) was dissolved in acetic anhydride by gentle warming. The resulting solution was stirred at room temperature overnight. The acetic anhydride was removed by evaporation under vacuum; the residue was dissolved in methylene chloride and morpholine (0.16 mL, 1.82 mmol) was added. The mixture was stirred overnight at room temperature and then concentrated under vacuum. The residue was dissolved in ethyl acetate, washed with 1 N hydrochloric acid solution, water and brine, dried over magnesium sulfate and concentrated to afford [(4-methoxybenzenesulfonyl)(2-morpholin-4-yl-2-oxoethyl)amino]acetic acid as an oil, 0.33 grams (54%).

To a solution of [(4-methoxybenzenesulfonyl)(2-morpholin-4-yl-2-oxoethyl)amino]acetic acid (0.33 grams, 0.89 mmol) in methylene chloride (10 mL) were added sequentially triethylamine (0.43 mL, 3.1 mmol), O-benzylhydroxylamine

- 50 -

hydrochloride (0.15 grams, 0.94 mmol) and (benzotriazol-1-yloxy)tris(dimethylamino)phosphonium hexafluorophosphate (0.43 grams, 0.97 mmol). The mixture was stirred at room temperature overnight and then diluted with ethyl acetate. The solution was washed sequentially with 0.5 N hydrochloric acid solution, water and brine, dried over magnesium sulfate and concentrated under vacuum. The residue was chromatographed on silica gel using ethyl acetate to afford N-benzyloxy-[(4-methoxybenzenesulfonyl)(2-morpholin-4-yl-2-oxoethyl)amino]acetamide as a white solid, 0.33 grams (78%).

To a solution of N-benzyloxy-[(4-methoxybenzenesulfonyl)(2-morpholin-4-yl-2-oxoethyl)amino]acetamide (0.33 grams, 0.69 mmol) in methanol (35 mL) was added 5% palladium on activated carbon (85 mg). The mixture was agitated under 2 atmospheres hydrogen in a Parr shaker for 1.5 hours. The catalyst was removed by filtration through nylon (pore size 0.45 μ m) and the solvent was evaporated. The residue was chromatographed on silica gel eluting with 5% methanol in methylene chloride to afford N-methoxy-[(4-methoxybenzenesulfonyl)(2-morpholin-4-yl-2-oxoethyl)amino]acetamide as a white solid, 65 mg (24%); ^1H NMR (CD_3OD): δ 7.82 (d, J = 9.0 Hz, 2H), 7.08 (d, J = 9.0 Hz, 2H), 4.24 (s, 2H), 3.88 (s, 3H), 3.84 (s, 2H), 3.68-3.64 (m, 4H), 3.58-3.50 (m, 4H); MS (thermospray): m/z 388 (M+1), 387 (M); HRMS calculated for $\text{C}_{16}\text{H}_{22}\text{N}_3\text{O}_7\text{S}$ (M+H): 388.1178, Found 338.1180.

The title compounds of Examples 15-16 were prepared by a method analogous to that described in Example 14 using [carboxymethyl(4-methoxybenzenesulfonyl)amino]acetic acid as the starting material which, after treatment with acetic anhydride, is coupled with the amine indicated.

- 51 -

EXAMPLE 15N-Hydroxy-[(4-methoxybenzenesulfonyl)(2-oxo-2-pyrrolidin-1-ylethyl)amino]acetamide

Coupled with pyrrolidine: ^1H NMR (DMSO-d_6):
5 δ 11.26 (br s, 1H), 8.89 (s, 1H), 7.81 (d, $J = 8.9$ Hz, 2H), 7.10 (d, $J = 8.9$ Hz, 2H), 4.09 (s, 2H), 3.85 (s, 3H), 3.74 (s, 2H), 3.45-3.25 (m, 4H), 1.93-1.72 (m, 4H); MS (thermospray): m/z 372 ($M+1$): Analysis calculated for $\text{C}_{15}\text{H}_{21}\text{N}_3\text{O}_6\text{S}$: C, 48.51; H, 5.70; N, 11.31. Found: C, 48.51; H, 5.82; N, 11.24.
10

EXAMPLE 162-[Dimethylcarbamoylmethyl(4-methoxybenzenesulfonyl)amino]-N-hydroxyacetamide

Coupled with dimethylamine: mp: 170°C .
15 (dec.); ^1H NMR (DMSO-d_6): δ 10.69 (br s, 1H), 8.88 (s, 1H), 7.91 (d, $J = 8.9$ Hz, 2H), 7.06 (d, $J = 8.9$ Hz, 2H), 4.19 (s, 2H), 3.85 (s, 3H), 3.73 (s, 2H), 2.94 (s, 3H), 2.84 (s, 3H); MS (thermospray): m/z 346 ($M+1$); Analysis calculated for $\text{C}_{13}\text{H}_{19}\text{N}_3\text{O}_6\text{S}$: C, 45.21; H, 5.55; N, 12.17. Found: C, 44.93; H, 5.61; N, 12.03.
20

EXAMPLE 172-(R)-2-[(2-Carbamoylethyl)(4-methoxybenzenesulfonyl)amino]-N-hydroxy-3-methylbutyramide

25 To a solution of 2-(R)-2-[(2-carboxylethyl)(4-methoxybenzenesulfonyl)amino]-3-methylbutyric acid benzyl ester (example 2) (900 mg., 2.0 mmol) in methylene chloride (10 mL) was added thionyl chloride (0.16 mL, 2.2 mmol). The reaction mixture was stirred
30 for 1.5 hours at room temperature and then concentrated in vacuo. After dissolving the residue in methylene chloride (10 mL), ammonia gas was bubbled through the solution for 0.5 minutes. The solution was stirred at room temperature overnight and was concentrated under
35 vacuum. Flash chromatography of the residue on silica gel eluting with 2% methanol in chloroform provided 2-(R)-2-[(2-carbamoylethyl)(4-

- 52 -

methoxybenzenesulfonyl)amino]-N-hydroxy-3-methylbutyric acid benzyl ester as a clear oil (275 mg, 31%).

To a solution of 2-(R)-2-[(2-carbamoylethyl)(4-methoxybenzenesulfonyl)amino]-N-hydroxy-3-methylbutyric acid benzyl ester (275 mg, 0.61 mmol) in ethanol (15 mL) was added 10% palladium on activated carbon (30 mg). The mixture was agitated under 3 atmospheres hydrogen in a Parr shaker for 5 hours. The catalyst was removed by filtration through diatomaceous earth and the solvent was evaporated leaving 2-(R)-2-[(2-carbamoylethyl)(4-methoxybenzenesulfonyl)amino]-N-hydroxy-3-methylbutyric acid as a white foam, 211 mg (96%).

To a solution of 2-(R)-2-[(2-carbamoylethyl)(4-methoxybenzenesulfonyl)amino]-N-hydroxy-3-methylbutyric acid (205 mg, 0.57 mmol) and 1-hydroxybenzotriazole hydrate (85 mg, 0.55 mmol) in dry dimethylformamide (5 mL) was added 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride (120 mg, 0.63 mmol). After stirring for 30 minutes, hydroxylamine hydrochloride (158 mg, 2.3 mmol) and then N-methylmorpholine (0.37 mL, 3.4 mmol) were added. The mixture was stirred at room temperature overnight and then diluted with ethyl acetate. The solution was washed with water and brine. The solution was then dried over magnesium sulfate and concentrated under vacuum to leave an oil which was chromatographed on silica gel eluting with 2% methanol in chloroform to afford 2-(R)-2-[(2-carbamoylethyl)(4-methoxybenzenesulfonyl)amino]-N-hydroxy-3-methylbutyramide as a white solid, 45 mg (21%); ^1H NMR (DMSO- d_6): δ 10.74 (br s, 1H), 8.91 (br s, 1H), 7.74 (d, J = 8.8 Hz, 2H), 7.33 (br s, 1H), 7.07 (d, J = 8.8 Hz, 2H), 6.79 (br s, 1H), 3.93-3.82 (m, 1H, overlapped), 3.83 (s, 3H), 3.64 (d, J = 10.7 Hz, 1H), 3.25-3.12 (m, 1H), 2.62-2.48 (m, 1H), 2.42-2.30 (m, 1H), 2.06-1.94 (m, 1H), 0.79 (d, J = 6.6 Hz, 3H), 0.76

- 53 -

(d, J = 6.6 Hz, 3H); MS (thermospray): m/z 374 (M+H).

EXAMPLE 18

2-(R)-2-[(2-tert-Butoxycarbonylethyl)(4-methoxybenzenesulfonyl)-amino]-N-hydroxy-3-methylbutyramide

A solution of N,N-dimethylformamide di-tert-butyl acetal (1.9 mL, 7.9 mmol) in toluene (15 mL) was added dropwise to a solution of 2-(R)-2-[(2-carboxyethyl(4-methoxybenzenesulfonyl)amino)-3-methylbutyric acid benzyl ester (example 2) 900 mg, 2.0 mmol) in toluene at 80°C. After heating for 2 hours at 80°C., the mixture was cooled and concentrated to leave an amber oil which was chromatographed on silica gel eluting with chloroform to afford 2-(R)-2-[(2-tert-butoxycarbonylethyl)(4-methoxybenzenesulfonyl)amino]-3-methylbutyric acid benzyl ester as an oil, 3.75 mg (37%).

To a solution of 2-(R)-2-[(2-tert-butoxycarbonylethyl)(4-methoxybenzenesulfonyl)amino]-3-methylbutyric acid benzyl ester (370 mg, 0.73 mmol) in ethanol (20 mL) was added 10% palladium on activated carbon (40 mg). The mixture was agitated under 3 atmospheres hydrogen in a Parr shaker for 5 hours. The catalyst was removed by filtration through diatomaceous earth and the solvent was evaporated leaving 2-(R)-2-[(2-tert-butoxycarbonylethyl)(4-methoxybenzenesulfonyl)amino]-3-methylbutyric acid as a white foam, 30 mg (100%).

To a solution of 2-(R)-2-[(2-tert-butoxycarbonylethyl)(4-methoxybenzenesulfonyl)amino]-3-methylbutyric acid (303 mg, 0.73 mmol) and 1-hydroxybenzotriazole hydrate (108 mg, 0.70 mmol) in dry dimethylformamide (10 mL) was added 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride (153 mg, 0.80 mmol). After stirring for 45 minutes, hydr xylamin hydrochloride (203 mg, 2.9 mmol) and then N-methylmorpholine (0.48 mL, 4.4 mmol) were added. The

- 54 -

mixtur was stirred at r om temperature overnight and then concentrated under vacuum. The residu was chromatographed on silica gel eluting with 2% methanol in chloroform and again with 10% ethyl acetate in
5 hexane to afford 2-(R)-2-[(2-tert-butoxycarbonylethyl)(4-methoxybenzenesulfonyl)amino]-N-hydroxy-3-methylbutyramide as a white foam, 135 mg (43%); ^1H NMR ($\text{DMSO}-d_6$): δ 10.77 (br s 1H), 7.74 (d, J = 8.9 Hz, 2H), 7.08 (d, J = 8.9 Hz, 2H), 3.93-3.82
10 (m, 1H, overlapped), 3.83 (s, 3H), 3.64 (d, J = 10.8 Hz, 1H), 3.26-3.14 (m, 1H), 2.70-2.60 (m, 1H), 2.50-2.38 (m, 1H), 2.04-1.91 (m, 1H), 1.38 (s, 9H), 0.78 (d, J = 6.5 Hz, 3H), 0.72 (d, J = 6.5 Hz, 3H); MS (thermospray): m/z 431 (M+H), 375, 314.

15

EXAMPLE 19


2-(R)-2-[2-Carboxyethyl)(4-methoxybenzenesulfonyl)amino]-N-hydroxy-3-methylbutyramide

To a solution of 2-(R)-2-[2-tert-butoxycarbonylethyl)(4-methoxybenzenesulfonyl)amino]-N-hydroxy-3-methylbutyramide (example 18) (100 mg, 0.23 mmol) in methylene chloride (1 mL) at 0°C. was added trifluoroacetic acid (1 mL). The mixture was allowed to warm to room temperature while stirring overnight.
20 After evaporation of the trifluoroacetic acid and methylene chloride, the residue was chromatographed on silica gel eluting with 5% methanol in chloroform. Concentration of the appropriate fractions afforded 2-(R)-2-[2-carboxyethyl)(4-methoxybenzenesulfonyl)amino]-
30 N-hydroxy-3-methylbutyramide as a white solid, 35 mg (41%). ^1H NMR ($\text{DMSO}-d_6$): δ 10.79 (br s, 1H), 8.97 (br s, 1H), 7.76 (d, J = 8.9 Hz, 2H), 7.09 (d, J = 8.9 Hz, 2H), 3.95-3.82 (m, 1H, overlapped), 3.84 (s, 3H), 3.66 (d, J = 10.8 Hz, 1H), 3.30-3.20 (m, 1H), 2.73-2.62 (m, 1H), 2.50-2.40 (m, 1H), 2.07-1.94 (m, 1H), 0.80 (d, J = 6.5 Hz, 3H), 0.74 (d, J = 6.5 Hz, 3H); MS
35 (thermospray): m/z 375 (M+H), 314.

- 55 -

C L A I M S

1. A compound of the formula




15 or the pharmaceutically acceptable salts thereof,
wherein

n is 1 to 6;

X is hydroxy, (C₁-C₆)alkoxy or NR¹R² wherein R¹ and R² are each independently selected from the group consisting of hydrogen, (C₁-C₆)alkyl, piperidyl, (C₁-C₆)alkylpiperidyl, (C₆-C₁₀)arylpiperidyl, (C₅-C₉)heteroarylpiperidyl, (C₆-C₁₀)aryl(C₁-C₆)alkylpiperidyl, (C₅-C₉)heteroaryl(C₁-C₆)alkylpiperidyl, (C₁-C₆)acylpiperidyl, (C₆-C₁₀)aryl, (C₅-C₉)heteroaryl, (C₆-C₁₀)aryl(C₁-C₆)alkyl, (C₅-C₉)heteroaryl(C₁-C₆)alkyl, (C₆-C₁₀)aryl(C₆-C₁₀)aryl, (C₆-C₁₀)aryl(C₆-C₁₀)aryl(C₁-C₆)alkyl, (C₃-C₆)cycloalkyl, (C₃-C₆)cycloalkyl(C₁-C₆)alkyl, R⁵(C₂-C₆)alkyl, (C₁-C₅)alkyl(CHR⁵)(C₁-C₆)alkyl wherein R⁵ is hydroxy, (C₁-C₆)acyloxy, (C₁-C₆)alkoxy, piperazino, (C₁-C₆)acylamino, (C₁-C₆)alkylthio, (C₆-C₁₀)arylthio, (C₁-C₆)alkylsulfinyl, (C₆-C₁₀)arylsulfinyl, (C₁-C₆)alkylsulfoxyl, (C₆-C₁₀)arylsulfoxyl, amino, (C₁-C₆)alkylamino, ((C₁-C₆)alkyl)₂amino, (C₁-C₆)acylpiperazino, (C₁-C₆)alkylpiperazino, (C₆-C₁₀)aryl(C₁-C₆)alkylpiperazino, (C₅-C₉)heteroaryl(C₁-C₆)alkylpiperazino, morpholino, thiomorpholino, piperidino or pyrrolidino; R⁶(C₁-C₆)alkyl, (C₁-C₅)alkyl(CHR⁶)(C₁-C₆)alkyl wherein R⁶ is piperidyl, (C₁-C₆)alkylpiperidyl, (C₆-C₁₀)arylpiperidyl, (C₆-C₁₀)aryl(C₁-C₆)alkylpiperidyl, (C₅-


- 56 -

- C_9)heteroarylpiperidyl or (C_5-C_9) heteroaryl (C_1-C_6) alkylpiperidyl; and $CH(R^7)COR^8$ wherein R^7 is hydrogen, (C_1-C_6) alkyl, (C_6-C_{10}) aryl (C_1-C_6) alkyl, (C_5-C_9) heteroaryl (C_1-C_6) alkyl, (C_1-C_6) alkylthio (C_1-C_6) alkyl, (C_6-C_{10}) arylthio (C_1-C_6) alkyl, (C_1-C_6) alkylsulfinyl (C_1-C_6) alkyl, (C_6-C_{10}) arylsulfinyl (C_1-C_6) alkyl, (C_1-C_6) alkylsulfonyl (C_1-C_6) alkyl, (C_6-C_{10}) arylsulfonyl (C_1-C_6) alkyl, hydroxy (C_1-C_6) alkyl, amino (C_1-C_6) alkyl, (C_1-C_6) alkylamino (C_1-C_6) alkyl, $((C_1-C_6)$ alkylamino) $_2(C_1-C_6)$ alkyl, $R^9R^{10}NCO(C_1-C_6)$ alkyl or $R^9OCO(C_1-C_6)$ alkyl wherein R^9 and R^{10} are each independently selected from the group consisting of hydrogen, (C_1-C_6) alkyl, (C_6-C_{10}) aryl (C_1-C_6) alkyl and (C_5-C_9) heteroaryl (C_1-C_6) alkyl; and R^8 is $R^{11}O$ or $R^{11}R^{12}N$ wherein R^{11} and R^{12} are each independently selected from the group consisting of hydrogen, (C_1-C_6) alkyl, (C_6-C_{10}) aryl (C_1-C_6) alkyl and (C_5-C_9) heteroaryl (C_1-C_6) alkyl; or R^1 and R^2 , or R^9 and R^{10} , or R^{11} and R^{12} may be taken together to form an azetidiny, pyrrolidinyl, morpholinyl, thiomorpholinyl, indolinyl, isoindolinyl, tetrahydroquinolinyl, tetrahydroisoquinolinyl, (C_1-C_6) acylpiperazinyl, (C_1-C_6) alkylpiperazinyl, (C_6-C_{10}) arylpiperazinyl, (C_5-C_9) heteroarylpiperazinyl or a bridged diazabicycloalkyl ring selected from the group consisting of




- 57 -

5



d



e

wherein r is 1, 2 or 3;

m is 1 or 2;

10 p is 0 or 1; and

Q is hydrogen, (C_1-C_3) alkyl or (C_1-C_6) acyl;

R^3 and R^4 are each independently selected

from the group consisting of hydrogen, (C_1-C_6) alkyl,

trifluoromethyl, trifluoromethyl (C_1-C_6) alkyl, $(C_1-$

15 $C_6)$ alkyl (difluoromethylene), $(C_1-$

$C_3)$ alkyl(difluoromethylene) (C_1-C_3) alkyl, (C_6-C_{10}) aryl,

(C_5-C_9) heteroaryl, (C_6-C_{10}) aryl (C_1-C_6) alkyl, $(C_5-$

$C_9)$ heteroaryl (C_1-C_6) alkyl, (C_6-C_{10}) aryl (C_6-C_{10}) aryl,

(C_6-C_{10}) aryl (C_6-C_{10}) aryl (C_1-C_6) alkyl, $(C_3-$

20 $C_6)$ cycloalkyl, (C_3-C_6) cycloalkyl (C_1-C_6) alkyl,

hydroxy (C_1-C_6) alkyl, (C_1-C_6) acyloxy (C_1-C_6) alkyl, $(C_1-$

$C_6)$ alkoxy (C_1-C_6) alkyl, piperazinyl (C_1-C_6) alkyl, $(C_1-$

$C_6)$ acylamino (C_1-C_6) alkyl, piperidyl, $(C_1-$

$C_6)$ alkylpiperidyl, (C_6-C_{10}) aryl (C_1-C_6) alkoxy $(C_1-$

25 $C_6)$ alkyl, (C_5-C_9) heteroaryl (C_1-C_6) alkoxy (C_1-C_6) alkyl,

(C_1-C_6) alkylthio (C_1-C_6) alkyl, (C_6-C_{10}) arylthio $(C_1-$

$C_6)$ alkyl, (C_1-C_6) alkylsulfinyl (C_1-C_6) alkyl, $(C_6-$

$C_{10})$ arylsulfinyl (C_1-C_6) alkyl, (C_1-C_6) alkylsulfonyl $(C_1-$

$C_6)$ alkyl, (C_6-C_{10}) arylsulfonyl (C_1-C_6) alkyl, amino $(C_1-$

30 $C_6)$ alkyl, (C_1-C_6) alkylamino (C_1-C_6) alkyl, $((C_1-$

$C_6)$ alkylamino) $_2(C_1-C_6)$ alkyl, $R^{13}CO(C_1-C_6)$ alkyl wherein

R^{13} is $R^{20}O$ or $R^{20}R^{21}N$ wherein R^{20} and R^{21} are each

independently selected from the group consisting of

hydrogen, (C_1-C_6) alkyl, (C_6-C_{10}) aryl (C_1-C_6) alkyl or


35 (C_5-C_9) heteroaryl (C_1-C_6) alkyl; $r R^{14}(C_1-C_6)$ alkyl

wherein R^{14} is (C_1-C_6) acylpiperazino, $(C_6-$

- 58 -

- C_{10})arylpiperazino, (C_5-C_9) heteroarylpiperazino, (C_1-C_6) alkylpiperazino, (C_6-C_{10}) aryl (C_1-C_6) alkylpiperazino, (C_5-C_9) heteroaryl (C_1-C_6) alkylpiperazino, morpholino, thiomorpholino, piperidino, pyrrolidino, piperidyl,
- 5 (C_1-C_6) alkylpiperidyl, (C_6-C_{10}) arylpiperidyl, (C_5-C_9) heteroarylpiperidyl, (C_6-C_{10}) aryl (C_1-C_6) alkylpiperidyl, (C_5-C_9) heteroaryl (C_1-C_6) alkylpiperidyl or (C_1-C_6) acylpiperidyl;
- 10 or R^3 and R^4 , or R^{20} and R^{21} may be taken together to form a (C_3-C_6) cycloalkyl, oxacyclohexyl, thiocyclohexyl, indanyl or tetralinyl ring or a group of the formula

15



- wherein R^{15} is hydrogen, (C_1-C_6) acyl, (C_1-C_6) alkyl,
- 20 (C_6-C_{10}) aryl (C_1-C_6) alkyl, (C_5-C_9) heteroaryl (C_1-C_6) alkyl or (C_1-C_6) alkylsulfonyl; and
- Ar is (C_6-C_{10}) aryl, (C_5-C_9) heteroaryl, (C_1-C_6) alkyl (C_6-C_{10}) aryl, (C_1-C_6) alkoxy (C_6-C_{10}) aryl, $((C_1-C_6)$ alkoxy) $_2$ (C_6-C_{10}) aryl, (C_6-C_{10}) aryloxy (C_6-C_{10}) aryl,
- 25 (C_5-C_9) heteroaryloxy (C_6-C_{10}) aryl, (C_1-C_6) alkyl (C_5-C_9) heteroaryl, (C_1-C_6) alkoxy (C_5-C_9) heteroaryl, $((C_1-C_6)$ alkoxy) $_2$ (C_5-C_9) heteroaryl, (C_6-C_{10}) aryloxy (C_5-C_9) heteroaryl, (C_5-C_9) heteroaryloxy (C_5-C_9) heteroaryl;
- with the proviso that when either R^1 or R^2 is
- 30 $CH(R^7)COR^8$ wherein R^7 and R^8 are as defined above, the other of R^1 or R^2 is hydrogen, (C_1-C_6) alkyl or benzyl.
2. A compound according to claim 1, wherein n is 2.
3. A compound according to claim 1, wherein Ar
- 35 is 4-methoxyphenyl or 4-phenoxyphenyl.
4. A compound according to claim 1, 2 or 3, wherein either R^3 or R^4 is not hydrogen.
5. A compound according to claim 1, wherein n is


- 59 -

1 and either R^1 or R^2 is hydrogen.

6. A compound according to claim 4, wherein X is hydroxy, Ar is 4-methoxyphenyl or 4-phenoxyphenyl.

7. A compound according to claim 4, wherein X is alkoxy, Ar is 4-methoxyphenyl or 4-phenoxyphenyl.

8. A compound according to claim 1, wherein Ar is 4-methoxyphenyl or 4-phenoxyphenyl and R^3 and R^4 are taken together to form (C₃-C₆)cycloalkanyl, oxacyclohexanyl, thiocyclohexanyl, indanyl or a group of the formula



wherein R^{15} is (C₁-C₆)acyl, (C₁-C₆)alkyl, (C₆-C₁₀)aryl(C₁-C₆)alkyl, (C₅-C₉)heteroaryl(C₁-C₆)alkyl or (C₁-C₆)alkylsulfonyl.

9. A compound according to claim 1, wherein n is 2, Ar is 4-methoxyphenyl or 4-phenoxyphenyl, R^1 and R^2 are taken together to form piperazinyl, (C₁-C₆)alkylpiperazinyl, (C₆-C₁₀)aryl piperazinyl or (C₅-C₉)heteroaryl(C₁-C₆)alkylpiperazinyl, and either R^3 or R^4 is not hydrogen or both R^3 and R^4 are not hydrogen.

10. A compound according to claim 1, wherein n is 2, Ar is 4-methoxyphenyl or 4-phenoxyphenyl, R^1 is hydrogen or (C₁-C₆)alkyl, R^2 is 2-pyridylmethyl, 3-pyridylmethyl or 4-pyridylmethyl, and either R^3 or R^4 is not hydrogen or both R^3 and R^4 are not hydrogen.

11. A compound according to claim 1, wherein n is 1, Ar is 4-methoxyphenyl or 4-phenoxyphenyl, R^1 is hydrogen, R^2 is 2-pyridylmethyl, 3-pyridylmethyl or 4-pyridylmethyl, and either R^3 or R^4 is not hydrogen or both R^3 and R^4 are not hydrogen.

12. A compound according to claim 2, wherein Ar is 4-methoxyphenyl, R^1 is hydrogen or (C₁-C₆)alkyl and R^2 is R^5 (C₂-C₆)alkyl wherein R^5 is morpholino,

- 60 -

thiomorpholino, piperidino, pyrrolidin, (C₁-C₆)acylpiperazino, (C₁-C₆)alkylpiperazino, (C₆-C₁₀)arylpiperazino, (C₅-C₉)heteroarylpiperazino, (C₆-C₁₀)aryl(C₁-C₆)alkylpiperazino or (C₅-C₉)heteroaryl(C₁-C₆)alkylpiperazino and either R³ or R⁴ is not hydrogen or both R³ and R⁴ are not hydrogen.

13. A compound according to claim 1, wherein n is 1, Ar is 4-methoxyphenyl or 4-phenoxyphenyl, R¹ is hydrogen, R² is R⁵(C₂-C₆)alkyl wherein R⁵ is morpholino, thiomorpholino, piperidino, pyrrolidino, (C₁-C₆)acylpiperazino, (C₁-C₆)alkylpiperazino, (C₆-C₁₀)arylpiperazino, (C₅-C₉)heteroarylpiperazino, (C₆-C₁₀)aryl(C₁-C₆)alkylpiperazino or (C₅-C₉)heteroaryl(C₁-C₆)alkylpiperazino and either R³ or R⁴ is not hydrogen or both R³ and R⁴ are not hydrogen.

14. A compound according to claim 1, wherein said compound is selected from:

- 2- (R) -N-Hydroxy-2- [(4-methoxybenzenesulfonyl) (3-morpholin-4-yl-3-oxopropyl) amino] -3-methylbutyramide;
- 2- (R) -2- [(2-Benzylcarbamoyl ethyl) (4-methoxybenzenesulfonyl) amino] -N-hydroxy-3-methylbutyramide;
- 2- (R) -N-Hydroxy-2- [(4-methoxybenzenesulfonyl) (2- [(pyridin-3-ylmethyl) carbamoyl] ethyl) amino] -3-methylbutyramide;
- 2- (R) -N-Hydroxy-2- [(4-methoxybenzenesulfonyl) [2- (methylpyridin-3-ylmethylcarbamoyl) ethyl] amino] -3-methylbutyramide;
- 4- (3- [1- (R) -1-Hydroxycarbamoyl-2-methylpropyl] (4-methoxybenzenesulfonyl) amino] propionyl) piperazine-1-carboxylic acid, tert-butyl ester;
- 2- (R) -N-Hydroxy-2- [(4-methoxybenzenesulfonyl) (3-oxo-3-piperazin-1-ylpropyl) amino] -3-methylbutyramide hydrochloride;
- 2- (R) -2- [(Benzylcarbamoylmethyl) (4-

- 61 -

- methoxybenzen sulfonyl)amin]N-hydroxy-3-methylbutyramide;
- 2- (R) -N-Hydroxy-2- [(4-methoxybenzenesulfonyl) - [(2-morpholin-4-yl)ethylcarbamoyl)methyl] amino) -3-methylbutyramide;
- 5 2- (R) -N-Hydroxy-2- [(4-methoxybenzenesulfonyl) ((pyridin-3-yl)methyl) carbamoyl)methyl] amino) -3-methylbutyramide;
- 2- (R) -3,3,3-Trifluoro-N-hydroxy-2- [(methoxybenzenesulfonyl) (3-morpholin-4-yl-3-oxopropyl) amino] propionamide;
- 10 2- (R) -N-Hydroxy-2- [(4-phenoxybenzenesulfonyl) [2- (methylpyridin-4-yl)methylcarbamoyl) ether] amino) -3-methylbutyramide;
- 15 4- [4-Methoxybenzenesulfonyl) (3-morpholin-4-yl-3-oxopropyl) amino] -1-methylpiperidene-4-carboxylic acid hydroxyamide;
- 2- (R) -N-Hydroxy-2- [(4-methoxybenzenesulfonyl) - [3- (4-methylpiperazin-1-yl) -3-oxopropyl] amino) -3-methylbutyramide;
- 20 2- (R) -2- [(2-Carboxyethyl) (4-methoxybenzenesulfonyl) amino] -N-hydroxy-3-methylbutyramide;
- [(2-Carboxyethyl) (3,4-dimethoxybenzenesulfonyl) amino] -N-hydroxy-acetamide;
- 25 2- (R) -2- [(2-Carbamoyl)ethyl) (4-methoxybenzenesulfonyl) amino] -N-hydroxy-3-methylbutyramide;
- 2- (R) , 3- (R) -3,N-Dihydroxy-2- [(4-methoxybenzenesulfonyl) (3-oxo-3-piperidin-1-yl)propyl) amino] -butyramide;
- 30 2- (R) -N-Hydroxy-2- [(4-methoxybenzenesulfonyl) [3- (methylpyridin-3-yl)methylcarbamoyl) propyl] amino) -3-methylbutyramide;
- 35 2- (R) -N-Hydroxy-2- [(4-methoxybenzenesulfonyl) [2- (methylcarboxymethylcarbam yl)ethyl] amino) -3-

- 62 -

methylbutyramide;

2-(R)-N-Hydroxy-2-((4-methoxybenzenesulfonyl)-[(1-methylpiperidin-4-ylcarbonyl)methyl]amino)-3-methylbutyramide;

5 2-(R)-2-Cyclohexyl-N-hydroxy-2-((4-methoxybenzenesulfonyl)-[3-(4-methylpiperazin-1-yl)-3-oxopropyl]amino)-acetamide; and

2-(R)-N-Hydroxy-2-[(methoxybenzenesulfonyl)(3-morpholin-4-yl-3-oxopropyl)amino]-4-(morpholin-4-yl)butyramide.

10 15. A pharmaceutical composition for (a) the treatment of a condition selected from the group consisting of arthritis, cancer, tissue ulceration, restenosis, periodontal disease, epidermolysis bullosa, scleritis and other diseases characterized by matrix metalloproteinase activity, AIDS, sepsis, septic shock
15 and other diseases involving the production of tumor necrosis factor (TNF) or (b) the inhibition of matrix metalloproteinases or the production of tumor necrosis
20 factor (TNF) in a mammal, including a human, comprising an amount of a compound of claim 1 or a pharmaceutically acceptable salt thereof, effective in such treatments and a pharmaceutically acceptable carrier.


25 16. A method for the inhibition of (a) matrix metalloproteinases or (b) the production of tumor necrosis factor (TNF) in a mammal, including a human, comprising administering to said mammal an effective amount of a compound of claim 1 or a pharmaceutically
30 acceptable salt thereof.

17. A method for treating a condition selected from the group consisting of arthritis, cancer, tissue ulceration, restenosis, periodontal disease, epidermolysis bullosa, scleritis and other diseases
35 characterized by matrix metalloproteinase activity, AIDS, sepsis, septic shock and other diseases involving the production of tumor necrosis factor (TNF) in a

- 63 -

mammal, including a human, comprising administering to said mammal an amount of a compound of claim 1 or a pharmaceutically acceptable salt thereof, effective in treating such a condition.

- 5 18. A method of preparing a compound of the formula



- 15 or the pharmaceutically acceptable salts thereof, wherein


n is 1 to 6;

- X is hydroxy, (C₁-C₆)alkoxy or NR¹R² wherein R¹ and R² are each independently selected from the group consisting of hydrogen, (C₁-C₆)alkyl, piperidyl, (C₁-C₆)alkylpiperidyl, (C₆-C₁₀)arylpiperidyl, (C₅-C₉)heteroarylpiperidyl, (C₆-C₁₀)aryl(C₁-C₆)alkylpiperidyl, (C₅-C₉)heteroaryl(C₁-C₆)alkylpiperidyl, (C₁-C₆)acylpiperidyl, (C₆-C₁₀)aryl, (C₅-C₉)heteroaryl, (C₆-C₁₀)aryl(C₁-C₆)alkyl, (C₅-C₉)heteroaryl(C₁-C₆)alkyl, (C₆-C₁₀)aryl(C₆-C₁₀)aryl, (C₆-C₁₀)aryl(C₆-C₁₀)aryl(C₁-C₆)alkyl, (C₃-C₆)cycloalkyl, (C₃-C₆)cycloalkyl(C₁-C₆)alkyl, R⁵(C₂-C₆)alkyl, (C₁-C₅)alkyl(CHR⁵)(C₁-C₆)alkyl wherein R⁵ is hydroxy, (C₁-C₆)acyloxy, (C₁-C₆)alkoxy, piperazino, (C₁-C₆)acylamino, (C₁-C₆)alkylthio, (C₆-C₁₀)arylthio, (C₁-C₆)alkylsulfinyl, (C₆-C₁₀)arylsulfinyl, (C₁-C₆)alkylsulfoxyl, (C₆-C₁₀)arylsulfoxyl, amino, (C₁-C₆)alkylamino, ((C₁-C₆)alkyl)₂amino, (C₁-C₆)acylpiperazino, (C₁-C₆)alkylpiperazino, (C₆-C₁₀)aryl(C₁-C₆)alkylpiperazino, (C₅-C₉)heteroaryl(C₁-
- 20
- 25
- 30
- 35

- 64 -

- C_6)alkylpiperazino, morpholino, thiomorpholino, piperidino or pyrrolidino; R^6 (C_1-C_6)alkyl, (C_1-C_5)alkyl(CHR^6) (C_1-C_6)alkyl wherein R^6 is piperidyl, (C_1-C_6)alkylpiperidyl, (C_6-C_{10})arylpiperidyl, (C_6-C_{10})aryl(C_1-C_6)alkylpiperidyl, (C_5-C_9)heteroarylpiperidyl or (C_5-C_9)heteroaryl(C_1-C_6)alkylpiperidyl; and $CH(R^7)COR^8$ wherein R^7 is hydrogen, (C_1-C_6)alkyl, (C_6-C_{10})aryl(C_1-C_6)alkyl, (C_5-C_9)heteroaryl(C_1-C_6)alkyl, (C_1-C_6)alkylthio(C_1-C_6)alkyl, (C_6-C_{10})arylthio(C_1-C_6)alkyl, (C_1-C_6)alkylsulfinyl(C_1-C_6)alkyl, (C_6-C_{10})arylsulfinyl(C_1-C_6)alkyl, (C_1-C_6)alkylsulfonyl(C_1-C_6)alkyl, (C_6-C_{10})arylsulfonyl(C_1-C_6)alkyl, hydroxy(C_1-C_6)alkyl, amino(C_1-C_6)alkyl, (C_1-C_6)alkylamino(C_1-C_6)alkyl, ($(C_1-C_6)alkylamino)_2(C_1-C_6)alkyl$, $R^9R^{10}NCO(C_1-C_6)alkyl$ or $R^9OCO(C_1-C_6)alkyl$ wherein R^9 and R^{10} are each independently selected from the group consisting of hydrogen, (C_1-C_6)alkyl, (C_6-C_{10})aryl(C_1-C_6)alkyl and (C_5-C_9)heteroaryl(C_1-C_6)alkyl; and R^8 is $R^{11}O$ or $R^{11}R^{12}N$ wherein R^{11} and R^{12} are each independently selected from the group consisting of hydrogen, (C_1-C_6)alkyl, (C_6-C_{10})aryl(C_1-C_6)alkyl and (C_5-C_9)heteroaryl(C_1-C_6)alkyl;
- or R^1 and R^2 , or R^9 and R^{10} , or R^{11} and R^{12}
- may be taken together to form an azetidiny, pyrrolidinyl, morpholinyl, thiomorpholinyl, indolinyl, isoindolinyl, tetrahydroquinolinyl, tetrahydroisoquinolinyl, (C_1-C_6)acylpiperazinyl, (C_1-C_6)alkylpiperazinyl, (C_6-C_{10})arylpiperazinyl, (C_5-C_9)heteroarylpiperazinyl or a bridged diazabicycloalkyl ring selected from the group consisting of

- 65 -



wherein r is 1, 2 or 3;

15

m is 1 or 2;

p is 0 or 1; and

Q is hydrogen, (C_1-C_3) alkyl or (C_1-C_6) acyl;

R^3 and R^4 are each independently selected

from the group consisting of hydrogen, (C_1-C_6) alkyl, trifluoromethyl, trifluoromethyl (C_1-C_6) alkyl, (C_1-C_6) alkyl (difluoromethylene), (C_1-C_3) alkyl (difluoromethylene) (C_1-C_3) alkyl, (C_6-C_{10}) aryl, (C_5-C_9) heteroaryl, (C_6-C_{10}) aryl (C_1-C_6) alkyl, (C_5-C_9) heteroaryl (C_1-C_6) alkyl, (C_6-C_{10}) aryl (C_6-C_{10}) aryl, (C_6-C_{10}) aryl (C_6-C_{10}) aryl (C_1-C_6) alkyl, (C_3-C_6) cycloalkyl, (C_3-C_6) cycloalkyl (C_1-C_6) alkyl, hydroxy (C_1-C_6) alkyl, (C_1-C_6) acyloxy (C_1-C_6) alkyl, (C_1-C_6) alkoxy (C_1-C_6) alkyl, piperazinyl (C_1-C_6) alkyl, (C_1-C_6) acylamino (C_1-C_6) alkyl, piperidyl, (C_1-C_6) alkylpiperidyl, (C_6-C_{10}) aryl (C_1-C_6) alkoxy (C_1-C_6) alkyl, C_5-C_9 heteroaryl (C_1-C_6) alkoxy (C_1-C_6) alkyl, (C_1-C_6) alkylthio (C_1-C_6) alkyl, (C_6-C_{10}) arylthio $(C_1-$

20


25

30

- 66 -

- C_6)alkyl, (C_1-C_6) alkylsulfinyl (C_1-C_6) alkyl, (C_6-C_{10}) arylsulfinyl (C_1-C_6) alkyl, (C_1-C_6) alkylsulfonyl (C_1-C_6) alkyl, (C_6-C_{10}) arylsulfonyl (C_1-C_6) alkyl, amino (C_1-C_6) alkyl, (C_1-C_6) alkylamino (C_1-C_6) alkyl, $((C_1-C_6)$ alkylamino) $_2$ (C_1-C_6) alkyl, $R^{13}CO(C_1-C_6)$ alkyl wherein R^{13} is $R^{20}O$ or $R^{20}R^{21}N$ wherein R^{20} and R^{21} are each independently selected from the group consisting of hydrogen, (C_1-C_6) alkyl, (C_6-C_{10}) aryl (C_1-C_6) alkyl or (C_5-C_9) heteroaryl (C_1-C_6) alkyl; or $R^{14}(C_1-C_6)$ alkyl wherein R^{14} is (C_1-C_6) acylpiperazino, (C_6-C_{10}) arylpiperazino, (C_5-C_9) heteroarylpiperazino, (C_1-C_6) alkylpiperazino, (C_6-C_{10}) aryl (C_1-C_6) alkylpiperazino, (C_5-C_9) heteroaryl (C_1-C_6) alkylpiperazino, morpholino, thiomorpholino, piperidino, pyrrolidino, piperidyl, (C_1-C_6) alkylpiperidyl, (C_6-C_{10}) arylpiperidyl, (C_5-C_9) heteroarylpiperidyl, (C_6-C_{10}) aryl (C_1-C_6) alkylpiperidyl, (C_5-C_9) heteroaryl (C_1-C_6) alkylpiperidyl or (C_1-C_6) acylpiperidyl; or R^3 and R^4 , or R^{20} and R^{21} may be taken together to form a (C_3-C_6) cycloalkyl, oxacyclohexyl, thiocyclohexyl, indanyl or tetralinyl ring or a group of the formula

25




- wherein R^{15} is hydrogen, (C_1-C_6) acyl, (C_1-C_6) alkyl, (C_6-C_{10}) aryl (C_1-C_6) alkyl, (C_5-C_9) heteroaryl (C_1-C_6) alkyl or (C_1-C_6) alkylsulfonyl; and
 Ar is (C_6-C_{10}) aryl, (C_5-C_9) heteroaryl, (C_1-C_6) alkyl (C_6-C_{10}) aryl, (C_1-C_6) alkoxy (C_6-C_{10}) aryl, $((C_1-C_6)$ alkoxy) $_2$ (C_6-C_{10}) aryl, (C_6-C_{10}) aryloxy (C_6-C_{10}) aryl, (C_5-C_9) heteroaryloxy (C_6-C_{10}) aryl, (C_1-C_6) alkyl (C_5-C_9) heteroaryl, (C_1-C_6) alkoxy (C_5-C_9) heteroaryl, $((C_1-C_6)$ alkoxy) $_2$ (C_5-C_9) heteroaryl, (C_6-C_{10}) aryloxy (C_5-C_9) heteroaryl, (C_5-C_9) heteroaryl xy (C_5-C_9) heteroaryl;

35

- 67 -

with the proviso that when either R^1 or R^2 is $CH(R^7)COR^8$ wherein R^7 and R^8 are as defined above, the other of R^1 or R^2 is hydrogen, (C_1-C_6) alkyl or benzyl; comprising reacting a compound of the formula



wherein n , X , R^3 , R^4 and Ar are as defined above with
 1- (3-dimethylaminopropyl) -3-ethylcarbodiimide, 1-
 15 hydroxybenztriazole and hydroxylamine.

INTERNATIONAL SEARCH REPORT

International Application No
PCT/US 96/02679

A. CLASSIFICATION OF SUBJECT MATTER
 IPC 6 C07C311/29 C07D295/18 C07D213/56 A61K31/535 A61K31/44
 A61K31/495 A61K31/40 A61K31/18

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)
 IPC 6 C07C C07D

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	EP,A,0 606 046 (CIBA-GEIGY AG) 13 July 1994 see the whole document ---	1-15,18
A	WO,A,90 05719 (BRITISH BIOTECHNOLOGY LTD) 31 May 1990 see the whole document -----	1-15,18

☐ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents:

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

- *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
- *A* document member of the same patent family

Date of the actual completion of the international search

15 May 1996

Date of mailing of the international search report

04.06.96

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
 NL - 2280 HV Rijswijk
 Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
 Fax: (+31-70) 340-3016

Authorized officer

Goetz, G

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US 96/02679

Patent document cited in search report	Publication date	Patent family member(s)	Publication date	
EP-A-606046	13-07-94	US-A-	5455258	03-10-95
		AU-B-	5265593	04-05-95
		CA-A-	2112779	07-07-94
		FI-A-	940012	07-07-94
		HU-A-	70536	30-10-95
		JP-A-	6256293	13-09-94
		NO-A-	940038	07-07-94
		NZ-A-	250517	26-10-95
		US-A-	5506242	09-04-96
		ZA-A-	9400048	11-08-94

WO-A-9005719	31-05-90	AU-B-	644064	02-12-93
		AU-B-	4800390	12-06-90
		CA-A-	2003718	23-05-90
		DE-D-	68914687	19-05-94
		DE-T-	68914687	08-09-94
		EP-A-	0446267	18-09-91
		ES-T-	2055409	16-08-94
		JP-T-	4502008	09-04-92
		NO-B-	177701	31-07-95
		US-A-	5310763	10-05-94
US-A-	5240958	31-08-93		
